

TABLE 3-5. Contraceptive use, number of previous abortions, and number of living children by age of patient, Oregon occurrence, 2017

Contraceptive used, previous abortions, and number of living children	Total	Age groups								
		<15	15-19	20-24	25-29	30-34	35-39	40-44	45+	N.S.
Total	8,506	15	817	2,374	2,375	1,601	975	323	24	2
None used	5,447	13	554	1,547	1,505	1,017	574	218	18	1
No previous abortion	3,463	13	494	1,142	877	546	273	109	8	1
One	1,129	–	51	290	359	237	126	58	8	–
Two	467	–	7	67	143	125	96	28	1	–
Three	177	–	1	17	63	48	36	12	–	–
Four or more	156	–	–	16	48	49	34	8	1	–
Pills used	933	–	102	291	303	133	89	14	1	–
No previous abortion	585	–	90	204	173	60	49	8	1	–
One	207	–	8	67	71	39	18	4	–	–
Two	80	–	1	9	39	18	11	2	–	–
Three	30	–	1	6	9	8	6	–	–	–
Four or more	24	–	1	3	9	7	4	–	–	–
Condoms used	1,065	2	78	262	268	222	181	47	4	1
No previous abortion	647	1	65	186	163	116	89	25	1	1
One	244	1	13	55	62	51	48	11	3	–
Two	99	–	–	11	26	31	25	6	–	–
Three	51	–	–	8	11	19	11	2	–	–
Four or more	22	–	–	1	6	4	8	3	–	–
Other contraceptive	821	–	48	194	232	203	111	32	1	–
No previous abortion	477	–	46	130	136	93	55	17	–	–
One	185	–	1	42	57	52	22	10	1	–
Two	88	–	1	12	24	28	20	3	–	–
Three	35	–	–	6	8	16	5	–	–	–
Four or more	29	–	–	1	5	12	9	2	–	–
Contraceptive use unknown ..	345	–	38	104	98	53	34	17	1	–
No previous abortion	220	–	31	70	55	29	23	11	1	–
One	80	–	6	27	22	16	6	3	–	–
Two	28	–	1	4	15	4	2	2	–	–
Three	4	–	–	1	2	1	–	–	–	–
Four or more	8	–	–	1	3	1	2	1	–	–
Number of living children										
No children ¹	4,181	15	724	1,486	1,115	567	216	52	5	1
Total with children	4,314	–	93	884	1,257	1,032	757	271	19	1
One	1,906	–	87	555	551	378	251	78	6	–
Two	1,460	–	6	270	425	374	280	100	5	–
Three	604	–	–	53	203	179	116	48	5	–
Four	221	–	–	3	57	58	74	26	2	1
Five or more	123	–	–	3	21	43	36	19	1	–

– Quantity is zero.

¹ Rows will not add to total due to some patients having an unknown number of children.

N.S. = Not stated.

NOTE: Contraceptive totals include abortions where the number of previous abortions is unknown. Multiple contraceptive methods may be reported for a single patient.