

TABLE 7-16.
PERINATAL DEATH RATES BY MOTHER'S RISK FACTORS,
OREGON RESIDENTS, BIRTH COHORT 1993-1995

RISK FACTOR	PERINATAL I ¹			PERINATAL II ²			NEONATAL ³	
	NO.	RATIO	RATE	NO.	RATIO	RATE	NO.	RATE
TOTAL	765	6.1	6.0	1,114	8.8	8.8	452	3.6
MARITAL STATUS								
MARRIED	506	5.6	5.6	734	8.2	8.1	280	3.1
UNMARRIED	259	7.2	7.2	380	10.5	10.5	172	4.8
AGE OF MOTHER								
15-19	101	6.4	6.4	156	9.9	9.8	65	4.1
20-24	192	5.8	5.8	277	8.3	8.3	124	3.7
25-29	213	6.1	6.1	285	8.1	8.1	121	3.5
30-34	137	5.0	5.0	204	7.5	7.4	82	3.0
35-39	84	7.1	7.0	138	11.6	11.5	41	3.4
40-44	31	12.8	12.7	45	18.6	18.3	13	5.4
45+	2	-	-	2	-	-	1	-
MOTHER'S ETHNICITY/RACE*								
WHITE (NON-HISPANIC)	612	5.9	5.9	893	8.6	8.6	371	3.6
HISPANIC (ALL RACES)	95	7.1	7.1	130	9.7	9.7	49	3.7
AFRICAN AMERICAN	25	9.3	9.3	42	15.7	15.5	19	7.1
INDIAN	8	4.7	4.7	12	7.1	7.1	5	3.0
OTHER NON-WHITE	23	5.2	5.2	30	6.8	6.8	8	1.8
MOTHER'S EDUCATION								
GRADE SCHOOL	62	8.6	8.6	78	10.8	10.8	42	5.8
HIGH SCHOOL +	647	5.5	5.5	952	8.2	8.1	374	3.2
START OF PRENATAL CARE	544	5.5	5.5	817	8.2	8.2	329	3.3
1ST TRIMESTER								
2ND TRIMESTER	140	6.6	6.6	190	9.0	9.0	77	3.7
3RD TRIMESTER	16	3.8	3.8	20	4.8	4.8	8	1.9
NO CARE	51	49.1	48.2	63	60.6	59.0	34	32.7
TOBACCO USE								
YES	163	7.1	7.1	243	10.6	10.5	91	4.0
NO	580	5.7	5.6	836	8.1	8.1	344	3.4
ALCOHOL USED								
YES	32	9.3	9.2	47	13.7	13.5	14	4.1
NO	701	5.8	5.8	1,020	8.5	8.4	412	3.4

*Unless specified, racial categories do not include Hispanic ethnicity.

NOTE: Because of unreported items, the sum of all categories may not equal the total.

- Quantity is zero, or rate is based on less than 5 events.

¹ Perinatal Definition I, generally used for international comparisons, includes fetal deaths at 28 weeks of gestation or more, and infant deaths of less than 7 days.

² Perinatal Definition II, includes fetal deaths at 20 weeks of gestation or more, and infant deaths of less than 28 days.

³ Neonatal deaths, includes infant deaths of less than 28 days.