

**TABLE 7-14. Perinatal Death Rates by County of Residence,
Oregon Residents, Birth Cohort 1997-1999**

| County of Residence | Perinatal I ¹ | | | Perinatal II ² | | | Neonatal ³ | |
|---------------------|--------------------------|------|-------|---------------------------|------|-------|-----------------------|------|
| | No. | Rate | Ratio | No. | Rate | Ratio | No. | Rate |
| Total | 790 | 5.9 | 5.9 | 1,108 | 8.2 | 8.3 | 485 | 3.6 |
| Baker | 4 | — | — | 6 | 11.2 | 11.3 | — | — |
| Benton | 16 | 6.5 | 6.5 | 20 | 8.1 | 8.1 | 11 | 4.5 |
| Clackamas | 57 | 4.6 | 4.6 | 81 | 6.5 | 6.6 | 38 | 3.1 |
| Clatsop | 4 | — | — | 5 | 4.2 | 4.2 | 1 | — |
| Columbia | 3 | — | — | 7 | 4.3 | 4.3 | 3 | — |
| Coos | 14 | 7.2 | 7.2 | 15 | 7.7 | 7.7 | 11 | 5.7 |
| Crook | 5 | 7.4 | 7.4 | 8 | 11.8 | 11.9 | 4 | — |
| Curry | 2 | — | — | 3 | — | — | — | — |
| Deschutes | 26 | 6.5 | 6.5 | 40 | 9.9 | 10.0 | 17 | 4.2 |
| Douglas | 30 | 8.8 | 8.8 | 34 | 10.0 | 10.0 | 19 | 5.6 |
| Gilliam | — | — | — | — | — | — | — | — |
| Grant | 4 | — | — | 6 | 22.7 | 23.0 | 3 | — |
| Harney | — | — | — | 1 | — | — | 1 | — |
| Hood River | 8 | 8.6 | 8.6 | 10 | 10.7 | 10.8 | 4 | — |
| Jackson | 34 | 5.4 | 5.4 | 54 | 8.5 | 8.6 | 24 | 3.8 |
| Jefferson | 6 | 6.6 | 6.6 | 8 | 8.7 | 8.8 | 3 | — |
| Josephine | 13 | 5.5 | 5.5 | 18 | 7.6 | 7.6 | 8 | 3.4 |
| Klamath | 20 | 8.0 | 8.1 | 26 | 10.4 | 10.5 | 12 | 4.8 |
| Lake | — | — | — | 1 | — | — | 1 | — |
| Lane | 77 | 6.9 | 6.9 | 100 | 9.0 | 9.0 | 47 | 4.2 |
| Lincoln | 13 | 10.0 | 10.0 | 22 | 16.8 | 17.0 | 7 | 5.4 |
| Linn | 21 | 4.8 | 4.8 | 33 | 7.5 | 7.6 | 15 | 3.4 |
| Malheur | 16 | 10.6 | 10.6 | 21 | 13.8 | 13.9 | 10 | 6.6 |
| Marion | 79 | 5.8 | 5.8 | 117 | 8.6 | 8.7 | 48 | 3.6 |
| Morrow | 2 | — | — | 3 | — | — | 3 | — |
| Multnomah | 160 | 5.8 | 5.8 | 220 | 8.0 | 8.0 | 82 | 3.0 |
| Polk | 9 | 4.3 | 4.4 | 15 | 7.2 | 7.3 | 6 | 2.9 |
| Sherman | — | — | — | — | — | — | — | — |
| Tillamook | 6 | 8.6 | 8.7 | 10 | 14.3 | 14.4 | 3 | — |
| Umatilla | 18 | 5.6 | 5.6 | 26 | 8.1 | 8.2 | 9 | 2.8 |
| Union | 5 | 5.6 | 5.6 | 8 | 8.9 | 9.0 | — | — |
| Wallowa | 2 | — | — | 3 | — | — | 2 | — |
| Wasco | 7 | 7.8 | 7.8 | 11 | 12.2 | 12.3 | 5 | 5.6 |
| Washington | 102 | 4.9 | 5.0 | 139 | 6.7 | 6.8 | 64 | 3.1 |
| Wheeler | — | — | — | — | — | — | — | — |
| Yamhill | 23 | 6.6 | 6.6 | 32 | 9.1 | 9.2 | 19 | 5.4 |
| Not Stated | 4 | — | — | 5 | — | — | 5 | — |

— Quantity is zero or rate/ratio is based on fewer than five occurrences.

¹ Perinatal Definition I includes fetal deaths at 28 weeks of gestation or more and infant deaths of less than 7 days.

² Perinatal Definition II includes fetal deaths at 20 weeks of gestation or more and infant deaths of less than 28 days.

³ Neonatal deaths include infant deaths of less than 28 days.

Perinatal I and Perinatal II ratios and Neonatal rate are per 1,000 births. Perinatal I rate includes all live births and fetal deaths at 28 weeks gestation or more. Perinatal II rate includes all live births and fetal deaths at 20 weeks of gestation or more.