TABLE 7-17. Neonatal, Postneonatal, and Infant Death Rates by Mother's Risk Factors, Oregon Residents, Birth Cohort 1999

Risk Factor	Neonatal ¹		Post- Neonatal ²		Infant ³	
	No.	Rate	No.	Rate	No.	Rate
Total	192	4.2	69	1.5	261	5.8
Marital Status						
Married	119	3.8	36	1.1	155	4.9
Unmarried	73	5.3	33	2.4	106	7.7
Mother's Age						
10-14	_	_	_	_	_	_
15-19	32	5.8	15	2.7	47	8.6
20-24	54	4.5	18	1.5	72	6.1
25-29	37	2.9	20	1.6	57	4.5
30-34	39	4.1	9	1.0	48	5.1
35-39	24	5.2	7	1.5	31	6.8
40-44	6	5.9	_	_	6	5.9
45+	_	_	_	_	_	_
Non-Hispanic						
White	137	4.0	52	1.5	189	5.5
African American	3	_	2	_	5	5.7
American Indian	4	_	2	_	6	9.2
Asian ⁴	9	4.3	_	_	9	4.3
Total Hispanic	37	5.4	12	1.7	49	7.1
Mother's Education						
8 th Grade or Less	9	3.3	6	2.2	15	5.5
Some High School	31	4.8	18	2.8	49	7.7
HS diploma/GED	66	4.5	27	1.8	93	6.3
More than High School	70	3.5	17	0.9	87	4.4
Start of Prenatal Care						
1 st Trimester	151	4.1	57	1.6	208	5.7
2 nd Trimester	28	4.0	10	1.4	38	5.5
3 rd Trimester	1	_	_	_	1	_
No Care	12	22.7	2	_	14	26.5
Tobacco Use						
Yes	36	5.5	18	2.8	54	8.3
No	141	3.7	49	1.3	190	5.0
Alcohol Use		•				2.0
Yes	3	_	4	_	7	8.5
No	172	4.0	63	1.5	235	5.4
Multiple Birth	.,_	1.0		1.5		О. т
Yes	17	14.7	4	_	21	18.2
No	173	3.9	65	1.5	238	5.4

Quantity is zero or rate is based on fewer than five occurrences.

NOTE: Because of unreported items, the sum of all categories may not equal the total. All rates per 1,000 live births.

¹ Neonatal deaths include infant deaths of less than 28 days.

² Postneonatal deaths occur from day 28 through 364 after birth.

³ Infant death is the death of a child prior to its first birthday.

⁴ Includes Chinese, Japanese, Filipino, and Other Asian & Pacific Islander.