

TABLE 7-9. Late Neonatal Deaths by Weeks of Gestation and Weight, Oregon Residents, Birth Cohort 2000

Birthweight (In Grams)	Total	Weeks of Gestation								
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+
Total	34	–	1	5	4	3	–	10	5	6
001-349	–	–	–	–	–	–	–	–	–	–
350-499	2	–	1	1	–	–	–	–	–	–
<500	2	–	1	1	–	–	–	–	–	–
500-749	3	–	–	2	1	–	–	–	–	–
750-999	4	–	–	2	2	–	–	–	–	–
1000-1249	1	–	–	–	1	–	–	–	–	–
1250-1499	1	–	–	–	–	1	–	–	–	–
1500-1999	1	–	–	–	–	–	–	–	–	1
2000-2499	2	–	–	–	–	–	–	1	1	–
<2500	14	–	1	5	4	1	–	1	1	1
2500-2999	7	–	–	–	–	2	–	1	1	3
3000-3499	5	–	–	–	–	–	–	4	1	–
3500-3999	5	–	–	–	–	–	–	3	2	–
4000-4499	3	–	–	–	–	–	–	1	–	2
4500-4999	–	–	–	–	–	–	–	–	–	–

– Quantity is zero.

Late neonatal death is defined as death at 7 to 27 days old.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used.

TABLE 7-10. Postneonatal Deaths by Weeks of Gestation and Weight, Oregon Residents, Birth Cohort 2000

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total	100	–	–	4	7	6	6	39	13	25	–
001-349	–	–	–	–	–	–	–	–	–	–	–
350-499	–	–	–	–	–	–	–	–	–	–	–
<500	–	–	–	–	–	–	–	–	–	–	–
500-749	–	–	–	–	–	–	–	–	–	–	–
750-999	5	–	–	3	2	–	–	–	–	–	–
1000-1249	4	–	–	1	2	1	–	–	–	–	–
1250-1499	2	–	–	–	2	–	–	–	–	–	–
1500-1999	4	–	–	–	1	1	1	1	–	–	–
2000-2499	12	–	–	–	–	1	3	6	–	2	–
<2500	27	–	–	4	7	3	4	7	–	2	–
2500-2999	21	–	–	–	–	2	1	11	2	5	–
3000-3499	29	–	–	–	–	1	1	12	5	10	–
3500-3999	20	–	–	–	–	–	–	8	4	8	–
4000-4499	1	–	–	–	–	–	–	–	1	–	–
4500-4999	2	–	–	–	–	–	–	1	1	–	–
Unknown	–	–	–	–	–	–	–	–	–	–	–

– Quantity is zero.

Postneonatal deaths occur from day 28 through 364 after birth.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used.