

**TABLE 7-10. Postneonatal Deaths by Weeks of Gestation and Weight,  
Oregon Residents, Birth Cohort 2002**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
<b>Total</b> .....	87	—	—	11	1	8	9	32	17	9	—
001-349 .....	—	—	—	—	—	—	—	—	—	—	—
350-499 .....	3	—	—	3	—	—	—	—	—	—	—
<500 .....	3	—	—	3	—	—	—	—	—	—	—
500-749 .....	2	—	—	2	—	—	—	—	—	—	—
750-999 .....	5	—	—	4	1	—	—	—	—	—	—
1000-1249 .....	3	—	—	2	—	1	—	—	—	—	—
1250-1499 .....	1	—	—	—	—	1	—	—	—	—	—
1500-1999 .....	4	—	—	—	—	2	1	1	—	—	—
2000-2499 .....	9	—	—	—	—	3	2	3	1	—	—
<2500 .....	27	—	—	11	1	7	3	4	1	—	—
2500-2999 .....	15	—	—	—	—	1	3	8	2	1	—
3000-3499 .....	28	—	—	—	—	—	2	17	8	1	—
3500-3999 .....	15	—	—	—	—	—	1	3	6	5	—
4000-4499 .....	2	—	—	—	—	—	—	—	—	2	—
4500-4999 .....	—	—	—	—	—	—	—	—	—	—	—
Unknown .....	—	—	—	—	—	—	—	—	—	—	—

— Quantity is zero.

Postneonatal deaths occur from day 28 through 364 after birth.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used.