

TABLE 6-11. Years of Potential Life Lost before Age 65 from the Leading Causes of Death, by Year, Oregon Residents, 1990-2004

Year	Total	Unintentional Injury	Cancer	Heart Disease ¹	Suicide	Perinatal Conditions	Congenital Anomalies	Alcohol-induced ²	Diabetes
1990	117,310	26,397	19,097	10,260	9,609	7,586	6,602	2,647	1,181
1991	113,112	23,842	19,215	11,005	9,801	6,291	6,710	2,582	1,388
1992	114,350	21,758	18,655	10,670	10,492	7,069	6,220	2,845	1,916
1993	123,280	25,797	19,747	12,169	9,772	5,391	7,125	3,334	1,594
1994	126,313	25,604	21,242	11,189	11,467	6,809	5,848	3,491	1,890
1995	128,177	28,912	20,505	12,226	12,029	4,932	5,394	3,856	1,811
1996	126,458	28,627	21,610	12,764	11,304	6,155	5,238	4,086	2,019
1997	120,508	27,322	21,233	12,748	10,937	6,596	5,867	3,783	2,036
1998	122,992	27,500	22,356	12,404	11,771	5,128	6,310	4,011	2,447
1999	117,350	21,710	21,254	13,390	9,807	7,276	6,523	3,142	2,441
2000.....	116,864	23,208	21,568	11,693	10,242	6,806	5,442	3,734	2,050
2001.....	118,229	22,052	22,574	11,589	10,566	7,276	5,651	4,484	2,422
2002.....	125,287	22,563	22,994	12,333	10,150	7,766	6,114	4,582	2,575
2003.....	126,196	25,182	21,504	12,676	10,716	7,441	5,225	5,522	3,376
2004.....	124,287	25,424	21,652	11,505	10,614	7,276	5,551	5,486	3,528

Year	Homicide ³	Cerebrovascular Disease	Undetermined External Cause	Chronic Lower Respiratory Disease	Acquired Immune Deficiency Syndrome	Viral Hepatitis	Sudden Infant Death Syndrome	Pneumonia and Influenza	Septicemia
1990	3,505	1,770	1,427	1,341	4,778	316	7,098	1,494	332
1991	4,152	1,801	1,112	1,309	5,796	288	5,484	900	113
1992	4,973	2,087	1,706	1,213	6,479	216	5,423	1,224	423
1993	4,475	2,399	1,746	1,424	7,884	475	5,873	1,469	302
1994	5,568	2,799	1,747	1,309	8,419	593	4,064	1,434	374
1995	5,139	2,052	2,021	1,509	8,214	678	4,906	901	205
1996	4,884	2,277	2,265	1,625	5,559	608	3,033	1,115	501
1997	4,081	2,432	1,413	1,660	2,286	663	2,323	1,313	185
1998	4,224	2,520	1,342	1,392	1,668	951	2,903	1,177	615
1999	3,724	2,226	1,596	1,720	1,700	620	1,679	768	975
2000.....	2,918	2,036	1,472	1,517	1,432	1,020	3,292	588	869
2001.....	2,938	2,583	1,910	1,485	1,417	923	1,872	968	684
2002.....	3,700	2,461	2,571	1,655	1,833	1,488	2,000	1,317	768
2003.....	2,662	2,504	2,628	1,927	1,776	1,189	1,484	1,092	658
2004.....	3,446	2,804	2,409	1,604	1,270	1,167	1,226	864	739

1 Includes alcoholic cardiomyopathy.

2 See Table 6-6, footnotes 35-36, for a list of included conditions and their ICD codes. Prior to 1999, figures do not include deaths due to alcohol poisoning.

3 Excludes legal intervention.