

**TABLE 7-9. Late Neonatal Deaths by Weeks of Gestation and Weight,  
Oregon Residents, Birth Cohort 2004**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
<b>Total</b> .....	43	—	—	12	9	2	2	11	5	2	—
001-349 .....	—	—	—	—	—	—	—	—	—	—	—
350-499 .....	—	—	—	—	—	—	—	—	—	—	—
<500 .....	—	—	—	—	—	—	—	—	—	—	—
500-749 .....	8	—	—	6	2	—	—	—	—	—	—
750-999 .....	7	—	—	6	1	—	—	—	—	—	—
1000-1249 .....	1	—	—	—	1	—	—	—	—	—	—
1250-1499 .....	2	—	—	—	2	—	—	—	—	—	—
1500-1999 .....	6	—	—	—	3	1	1	—	—	1	—
2000-2499 .....	5	—	—	—	—	—	1	2	2	—	—
<2500 .....	29	—	—	12	9	1	2	2	2	1	—
2500-2999 .....	5	—	—	—	—	—	—	5	—	—	—
3000-3499 .....	4	—	—	—	—	—	—	3	1	—	—
3500-3999 .....	3	—	—	—	—	1	—	—	2	—	—
4000-4499 .....	1	—	—	—	—	—	—	1	—	—	—
4500-4999 .....	1	—	—	—	—	—	—	—	—	1	—
N.S. ....	—	—	—	—	—	—	—	—	—	—	—

— Quantity is zero.

Late neonatal death is defined as death at 7 to 27 days old.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used.