

TABLE 6-54. Age-Adjusted Death Rates for Selected Causes of Death, by Year, Oregon Residents and the US, 1980-2005

Year	Total			Cancer			Heart Disease		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1980.....	965.3	1,038.7	-7.1	198.5	209.8	-5.4	350.4	405.9	-13.7
1985.....	949.2	987.8	-3.9	211.4	213.2	-0.8	334.0	369.3	-9.6
1990.....	866.0	938.0	-7.7	210.3	217.9	-3.5	255.5	317.0	-19.4
1995.....	882.3	909.5	-3.0	214.2	211.7	1.2	232.4	289.0	-19.6
1996.....	881.9	893.7	-1.3	208.8	208.6	0.1	230.6	281.4	-18.1
1997.....	864.0	877.5	-1.5	205.7	205.3	0.2	221.8	273.5	-18.9
1998.....	862.9	870.1	-0.8	207.9	202.5	2.7	210.7	267.2	-21.2
1999.....	845.3	875.6	-3.5	199.2	200.8	-0.8	208.0	266.4	-21.9
2000.....	826.9	869.0	-4.8	197.6	199.6	-1.0	197.5	257.6	-23.3
2001.....	835.9	854.5	-2.2	198.7	196.0	1.4	195.2	247.8	-18.9
2002.....	855.0	845.3	1.1	200.9	193.5	3.8	198.0	240.8	-14.8
2003.....	838.4	832.7	0.7	198.3	190.1	4.3	189.5	232.3	-12.7
2004.....	814.8	800.8	1.7	196.7	185.8	5.9	179.2	217.0	-17.4
2005*.....	791.4	798.8	-0.9	189.4	183.8	3.0	169.5	210.3	-19.4

Year	Cerebrovascular Disease			Chronic Lower Resp. Disease			Unintentional Injuries		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1980.....	101.0	101.2	-0.2	35.3	29.5	19.8	53.4	48.9	9.2
1985.....	86.4	80.4	7.4	43.4	35.9	20.8	48.0	40.9	17.4
1990.....	69.4	68.8	0.9	45.3	38.7	17.0	38.9	38.4	1.2
1995.....	81.6	66.5	22.8	46.6	41.8	11.4	41.3	36.4	13.5
1996.....	83.7	65.7	27.3	52.5	42.4	23.9	40.6	36.6	11.0
1997.....	80.8	64.3	25.7	50.8	42.8	18.7	39.9	36.4	9.7
1998.....	80.7	62.4	29.4	49.6	43.5	14.0	40.8	36.7	11.2
1999.....	80.3	61.6	30.4	50.4	45.4	11.0	33.9	34.8	-2.6
2000.....	70.8	60.9	16.3	47.8	44.2	8.1	34.6	34.5	0.3
2001.....	71.4	57.9	23.3	48.7	43.7	11.4	35.4	35.1	0.9
2002.....	71.7	56.2	27.6	50.9	43.5	17.0	38.4	36.3	5.8
2003.....	68.5	53.5	28.0	49.8	43.3	15.0	38.3	36.7	4.4
2004.....	61.9	50.0	23.8	48.1	41.1	17.0	38.8	37.2	4.3
2005*.....	57.3	46.6	23.0	47.8	43.2	10.6	37.6	38.1	-1.3

Note: US age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database, except for the most recent year which is a preliminary estimate from the National Center for Health Statistics. Unlike the data shown in Table 6-52, all Oregon data are from state mortality files, except for 1980 and 1985 which are from the CDC's online database. Consequently, the rates and percentage differences shown here will vary from those in Table 6-52 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. National comparability ratios have been applied to 1980-1998 rates to control for coding changes resulting from the implementation of ICD-10 when it supplanted ICD-9 in 1999. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's disease and Alzheimer's dementia). Some differences between Oregon and US rates (e.g., alcohol-induced deaths) result, at least in part, from the state's query program (when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information).

* National rates are preliminary.

TABLE 6-54. Age-Adjusted Death Rates for Selected Causes of Death, by Year, Oregon Residents and the US, 1980-2005 — Continued

Year	Alzheimer's Disease			Diabetes Mellitus			Flu & Pneumonia		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1980.....	1.7	1.1	53.6	13.5	18.4	-26.8	24.8	21.9	13.3
1985.....	10.3	6.5	58.9	12.8	17.7	-27.8	25.2	24.0	4.8
1990.....	15.2	10.1	50.2	16.8	21.1	-20.4	23.7	25.6	-7.6
1995.....	19.8	13.3	49.1	22.4	23.6	-5.2	19.7	23.3	-15.4
1996.....	20.6	13.4	53.3	23.0	24.3	-5.2	20.1	22.9	-12.3
1997.....	19.8	13.8	44.0	24.9	24.2	3.1	19.0	23.2	-18.1
1998.....	19.0	13.6	39.7	26.0	24.6	5.9	20.7	24.1	-14.2
1999.....	24.7	16.5	49.7	24.7	25.0	-1.2	19.5	23.5	-17.0
2000.....	24.8	18.0	37.8	23.8	25.0	-4.8	17.5	23.7	-26.2
2001.....	28.1	19.1	47.1	28.8	25.3	13.8	15.7	22.0	-28.6
2002.....	30.3	20.2	50.0	28.6	25.4	12.6	17.9	22.6	-20.8
2003.....	30.6	21.4	43.0	28.1	25.3	11.1	17.0	22.0	-22.7
2004.....	33.4	21.8	53.2	29.0	24.5	18.4	14.7	19.8	-25.8
2005*.....	30.4	22.9	32.8	29.3	25.2	16.3	15.1	20.3	-25.6

Year	Suicide			Alcohol-induced			Hypertension		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1980.....	15.1	12.2	23.5	11.9	9.8	21.4	2.9	4.8	-39.6
1985.....	16.0	12.5	27.7	12.0	8.3	44.6	3.7	4.2	-12.8
1990.....	15.8	12.5	26.1	11.8	8.9	32.6	4.9	4.6	7.1
1995.....	16.8	11.8	42.1	11.3	8.4	34.5	6.7	5.6	20.1
1996.....	16.7	11.5	44.9	12.8	8.2	56.1	6.5	5.7	14.2
1997.....	16.7	11.2	48.8	11.5	7.9	45.6	7.7	5.8	32.7
1998.....	17.2	11.1	54.6	11.0	7.8	41.0	6.6	6.0	9.5
1999.....	14.9	10.5	41.9	8.9	7.1	25.4	7.0	6.2	12.9
2000.....	14.3	10.4	37.5	10.8	7.0	54.3	6.2	6.6	-6.1
2001.....	14.9	10.7	39.3	12.2	7.0	74.3	8.6	6.8	26.5
2002.....	14.5	10.9	33.0	12.3	6.9	78.3	9.6	7.0	37.1
2003.....	16.3	10.7	52.3	14.2	7.0	102.9	9.3	7.4	25.7
2004.....	15.2	10.9	39.4	13.8	7.0	97.1	9.5	7.7	23.4
2005*.....	14.9	10.6	40.6	13.7	6.9	98.6	10.6	8.0	32.5

Note: US age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database, except for the most recent year which is a preliminary estimate from the National Center for Health Statistics. Unlike the data shown in Table 6-52, all Oregon data are from state mortality files, except for 1980 and 1985 which are from the CDC's online database. Consequently, the rates and percentage differences shown here will vary from those in Table 6-52 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. National comparability ratios have been applied to 1980-1998 rates to control for coding changes resulting from the implementation of ICD-10 when it supplanted ICD-9 in 1999. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's disease and Alzheimer's dementia). Some differences between Oregon and US rates (e.g., alcohol-induced deaths) result, at least in part, from the state's query program (when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information).

* National rates are preliminary.

TABLE 6-54. Age-Adjusted Death Rates for Selected Causes of Death, by Year, Oregon Residents and the US, 1980-2005 — Continued

Year	Parkinson's Disease			Arteriosclerosis			Amyotrophic Lateral Sclerosis		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1980.....	3.4	2.1	61.9	23.0	17.4	32.3	2.1	1.3	61.5
1985.....	3.9	2.6	50.0	17.5	12.2	43.8	1.7	1.4	21.4
1990.....	5.0	3.3	51.5	11.3	8.2	37.6	1.8	1.5	20.0
1995.....	7.2	4.3	67.4	9.0	6.6	37.0	1.9	1.6	18.8
1996.....	7.2	4.6	56.5	7.5	6.4	17.6	2.0	1.6	25.0
1997.....	6.4	4.7	36.2	6.9	6.0	15.2	2.3	1.6	43.8
1998.....	8.0	4.9	63.3	6.5	5.6	16.0	2.2	1.6	37.5
1999.....	7.3	5.4	35.2	5.6	5.5	1.8	2.2	1.9	15.8
2000.....	7.7	5.7	35.1	6.4	5.2	23.1	2.7	2.0	35.0
2001.....	8.0	5.9	35.6	5.3	5.0	6.0	2.6	1.9	35.8
2002.....	8.3	5.9	40.7	5.7	4.8	18.8	3.0	2.0	47.5
2003.....	8.4	6.2	35.5	5.5	4.4	25.0	3.1	2.0	55.5
2004.....	8.6	6.1	41.0	4.6	3.9	17.9	2.9	1.9	54.7
2005*.....	7.7	6.4	20.3	4.8	3.8	26.3	2.8	NA	NA

Year	Homicide			Viral Hepatitis			HIV/AIDS		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1980.....	5.1	10.3	-50.5	0.3	0.3	0.0	NA	NA	NA
1985.....	4.4	7.9	-44.3	0.4	0.3	33.3	NA	NA	NA
1990.....	3.7	9.4	-60.6	0.8	0.5	59.8	7.2	10.6	-32.1
1995.....	4.9	8.3	-41.0	1.5	0.9	61.4	11.5	16.8	-31.4
1996.....	4.5	7.5	-40.0	1.1	1.0	9.9	7.6	11.9	-36.1
1997.....	3.9	7.0	-44.3	1.4	1.1	30.5	3.2	6.2	-48.1
1998.....	4.1	6.4	-35.9	1.6	1.3	24.3	2.3	4.9	-52.8
1999.....	3.3	5.9	-44.1	1.3	1.8	-27.8	2.2	5.3	-58.5
2000.....	2.7	5.8	-53.4	2.2	1.9	15.8	1.8	5.2	-65.4
2001.....	3.1	6.0	-48.3	2.5	2.0	25.0	1.9	5.0	-62.0
2002.....	3.1	6.0	-48.3	3.5	2.0	75.0	2.5	4.9	-49.0
2003.....	2.5	6.0	-58.3	2.6	1.8	44.4	2.5	4.7	-46.8
2004.....	3.1	5.8	-46.6	2.9	1.8	61.1	1.8	4.5	-60.0
2005*.....	2.9	5.9	-50.8	2.3	1.8	27.8	1.5	4.2	-64.3

Note: US age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database, except for the most recent year which is a preliminary estimate from the National Center for Health Statistics. Unlike the data shown in Table 6-52, all Oregon data are from state mortality files, except for 1980 and 1985 which are from the CDC's online database. Consequently, the rates and percentage differences shown here will vary from those in Table 6-52 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. National comparability ratios have been applied to 1980-1998 rates to control for coding changes resulting from the implementation of ICD-10 when it supplanted ICD-9 in 1999. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's disease and Alzheimer's dementia). Some differences between Oregon and US rates (e.g., alcohol-induced deaths) result, at least in part, from the state's query program (when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information).

* National rates are preliminary.