

**TABLE 7-9. Late Neonatal Deaths¹ by Weeks of Gestation and Weight
Oregon Residents, Birth Cohort 2007**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total	35	–	1	8	5	5	3	12	1	–	–
001-349	–	–	–	–	–	–	–	–	–	–	–
350-499	1	–	–	1	–	–	–	–	–	–	–
<500	1	–	–	1	–	–	–	–	–	–	–
500-749	4	–	1	3	–	–	–	–	–	–	–
750-999	4	–	–	3	1	–	–	–	–	–	–
1000-1249	3	–	–	1	2	–	–	–	–	–	–
1250-1499	1	–	–	–	–	1	–	–	–	–	–
1500-1999	5	–	–	–	1	2	1	1	–	–	–
2000-2499	5	–	–	–	1	2	–	2	–	–	–
<2500	23	–	1	8	5	5	1	3	–	–	–
2500-2999	3	–	–	–	–	–	–	3	–	–	–
3000-3499	5	–	–	–	–	–	1	3	1	–	–
3500-3999	4	–	–	–	–	–	1	3	–	–	–
4000-4499	–	–	–	–	–	–	–	–	–	–	–
4500+	–	–	–	–	–	–	–	–	–	–	–
Unknown	–	–	–	–	–	–	–	–	–	–	–

– Quantity is zero.

¹ Late neonatal deaths occur from day 7 through 27 after birth.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.