

**TABLE 7-10. Postneonatal Deaths¹ by Weeks of Gestation and Weight
Oregon Residents, Birth Cohort 2007**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total	85	–	–	5	5	13	3	37	14	7	1
001-349	–	–	–	–	–	–	–	–	–	–	–
350-499	–	–	–	–	–	–	–	–	–	–	–
<500	–	–	–	–	–	–	–	–	–	–	–
500-749	5	–	–	5	–	–	–	–	–	–	–
750-999	2	–	–	–	2	–	–	–	–	–	–
1000-1249	1	–	–	–	1	–	–	–	–	–	–
1250-1499	3	–	–	–	2	1	–	–	–	–	–
1500-1999	7	–	–	–	–	6	–	1	–	–	–
2000-2499	9	–	–	–	–	5	–	3	1	–	–
<2500	27	–	–	5	5	12	–	4	1	–	–
2500-2999	19	–	–	–	–	–	3	12	2	2	–
3000-3499	18	–	–	–	–	1	–	13	3	1	–
3500-3999	16	–	–	–	–	–	–	6	7	3	–
4000-4499	4	–	–	–	–	–	–	2	1	1	–
4500+	–	–	–	–	–	–	–	–	–	–	–
Unknown	1	–	–	–	–	–	–	–	–	–	1

– Quantity is zero.

¹ Postneonatal deaths occur from day 28 through 364 after birth.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.