

TABLE 6-57. Age-Adjusted Death Rates for Selected Causes of Death, by Year, Oregon Residents and the US , 1985-2008

Year	Total			Cancer			Heart Disease		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1985	949.2	987.8	-3.9	211.4	213.2	-0.8	334.0	369.3	-9.6
1990	866.0	938.0	-7.7	210.3	217.9	-3.5	255.5	317.0	-19.4
1995	882.3	909.5	-3.0	214.2	211.7	1.2	232.4	289.0	-19.6
1996	881.9	893.7	-1.3	208.8	208.6	0.1	230.6	281.4	-18.1
1997	864.0	877.5	-1.5	205.7	205.3	0.2	221.8	273.5	-18.9
1998	862.9	870.1	-0.8	207.9	202.5	2.7	210.7	267.2	-21.2
1999	845.3	875.6	-3.5	199.2	200.8	-0.8	208.0	266.4	-21.9
2000	826.9	869.0	-4.8	197.6	199.6	-1.0	197.5	257.6	-23.3
2001	835.9	854.5	-2.2	198.7	196.0	1.4	195.2	247.8	-18.9
2002	855.0	845.3	1.1	200.9	193.5	3.8	198.0	240.8	-14.8
2003	838.4	832.7	0.7	198.3	190.1	4.3	189.5	232.3	-12.7
2004	814.8	800.8	1.7	196.7	185.8	5.9	179.2	217.0	-17.4
2005	791.4	798.8	-0.9	189.4	183.8	3.0	169.5	211.1	-19.7
2006	784.5	776.5	1.0	185.7	180.7	2.8	162.6	200.2	-18.8
2007	771.6	760.2	1.5	184.7	178.4	3.5	159.7	190.9	-16.3
2008	772.8	758.3	1.9	182.8	175.3	4.3	154.5	186.5	-17.2

Year	Cerebrovascular Disease			Chronic Lower Resp. Disease			Unintentional Injuries		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1985	86.4	80.4	7.5	43.4	35.9	20.9	48.0	40.9	17.4
1990	69.4	68.8	0.9	45.3	38.7	17.1	38.9	38.4	1.3
1995	81.6	66.5	22.7	46.6	41.8	11.5	41.3	36.4	13.5
1996	83.7	65.7	27.4	52.5	42.4	23.8	40.6	36.6	10.9
1997	80.8	64.3	25.7	50.8	42.8	18.7	39.9	36.4	9.6
1998	80.7	62.4	29.3	49.6	43.5	14.0	40.8	36.7	11.2
1999	80.3	61.6	30.4	50.4	45.4	11.0	33.9	34.8	-2.6
2000	70.8	60.9	16.3	47.8	44.2	8.1	34.6	34.5	0.3
2001	71.4	57.9	23.3	48.7	43.7	11.4	35.4	35.1	0.9
2002	71.7	56.2	27.6	50.9	43.5	17.0	38.4	36.3	5.8
2003	68.5	53.5	28.0	49.8	43.3	15.0	38.3	36.7	4.4
2004	61.9	50.0	23.8	48.1	41.1	17.0	38.8	37.2	4.3
2005	57.3	46.6	23.0	47.8	43.2	10.6	37.6	39.1	-3.8
2006	48.8	43.6	11.9	46.8	40.5	15.6	40.7	39.8	2.3
2007	44.5	42.2	5.5	47.5	40.8	16.4	41.7	40.0	4.3
2008	45.6	40.7	12.0	48.2	44.0	9.5	42.4	38.8	9.3

Note: US age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database. Unlike the data shown in Table 6-55, all Oregon data are from state mortality files, except for 1980 and 1985 which are from the CDC's online database. Consequently, the rates and percentage differences shown here will vary from those in Table 6-55 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. National comparability ratios have been applied to 1980-1998 rates to control for coding changes resulting from the implementation of ICD-10 when it supplanted ICD-9 in 1999. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's Disease and Alzheimer's dementia). Some differences between Oregon and US rates (e.g., alcohol-induced deaths) result, at least in part, from the state's query program (when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information).

TABLE 6-57. Age-Adjusted Death Rates for Selected Causes of Death, by Year, Oregon Residents and the US , 1985-2008 — Continued

Year	Alzheimer's Disease			Diabetes Mellitus			Suicide		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1985	10.3	6.5	58.5	12.8	17.7	-27.8	16.0	12.5	28.0
1990	15.2	10.1	50.5	16.8	21.1	-20.4	15.8	12.5	26.4
1995	19.8	13.3	48.9	22.4	23.6	-5.2	16.8	11.8	42.4
1996	20.6	13.4	53.7	23.0	24.3	-5.2	16.7	11.5	45.2
1997	19.8	13.8	43.5	24.9	24.2	3.1	16.7	11.2	49.1
1998	19.0	13.6	39.7	26.0	24.6	5.9	17.2	11.1	55.0
1999	24.7	16.5	49.7	24.7	25.0	-1.2	14.9	10.5	41.9
2000	24.8	18.0	37.8	23.8	25.0	-4.8	14.3	10.4	37.5
2001	28.1	19.1	47.1	28.8	25.3	13.8	14.9	10.7	39.3
2002	30.3	20.2	50.0	28.6	25.4	12.6	14.5	10.9	33.0
2003	30.6	21.4	43.0	28.1	25.3	11.1	16.3	10.7	52.3
2004	33.4	21.8	53.2	29.0	24.5	18.4	15.2	10.9	39.4
2005	30.4	22.9	32.8	29.3	24.6	19.1	14.9	10.9	36.7
2006	29.5	22.6	30.5	28.9	23.3	24.0	15.1	10.9	38.5
2007	28.0	22.7	23.3	27.9	22.5	24.0	15.6	11.3	38.1
2008	30.5	24.4	25.0	24.8	21.8	13.8	14.7	11.6	26.7

Year	Flu & Pneumonia			Alcohol-Induced			Hypertension		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1985	25.2	24.0	5.0	12.0	8.3	44.6	3.7	4.2	-12.8
1990	23.7	25.6	-7.6	11.8	8.9	32.6	4.9	4.6	6.5
1995	19.7	23.3	-15.4	11.3	8.4	34.5	6.7	5.6	19.6
1996	20.1	22.9	-12.3	12.8	8.2	56.1	6.5	5.7	14.0
1997	19.0	23.2	-18.1	11.5	7.9	45.6	7.7	5.8	32.8
1998	20.7	24.1	-14.2	11.0	7.8	41.0	6.6	6.0	10.0
1999	19.5	23.5	-17.0	8.9	7.1	25.4	7.0	6.2	12.9
2000	17.5	23.7	-26.2	10.8	7.0	54.3	6.2	6.6	-6.1
2001	15.7	22.0	-28.6	12.2	7.0	74.3	8.6	6.8	26.5
2002	17.9	22.6	-20.8	12.3	6.9	78.3	9.6	7.0	37.1
2003	17.0	22.0	-22.7	14.2	7.0	102.9	9.3	7.4	25.7
2004	14.7	19.8	-25.8	13.8	7.0	97.1	9.5	7.7	23.4
2005	15.1	20.3	-25.6	13.7	7.0	95.7	10.6	8.0	32.5
2006	12.8	17.8	-28.1	11.7	7.0	67.1	8.9	7.5	18.7
2007	11.4	16.2	-29.6	13.1	7.7	70.1	8.6	7.4	16.2
2008	12.3	16.9	-27.2	12.9	7.4	74.3	9.5	7.7	23.4

Note: US age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database. Unlike the data shown in Table 6-55, all Oregon data are from state mortality files, except for 1980 and 1985 which are from the CDC's online database. Consequently, the rates and percentage differences shown here will vary from those in Table 6-55 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. National comparability ratios have been applied to 1980-1998 rates to control for coding changes resulting from the implementation of ICD-10 when it supplanted ICD-9 in 1999. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's Disease and Alzheimer's dementia). Some differences between Oregon and US rates (e.g., alcohol-induced deaths) result, at least in part, from the state's query program (when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information).

TABLE 6-57. Age-Adjusted Death Rates for Selected Causes of Death, by Year, Oregon Residents and the US , 1985-2008 — Continued

Year	Parkinson's Disease			Homicide			Amyotrophic Lateral Sclerosis		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1985	3.9	2.6	50.0	4.4	7.9	-44.3	1.7	1.4	21.4
1990	5.0	3.3	51.5	3.7	9.4	-60.6	1.8	1.5	20.0
1995	7.2	4.3	67.4	4.9	8.3	-41.0	1.9	1.6	18.8
1996	7.2	4.6	56.5	4.5	7.5	-40.0	2.0	1.6	25.0
1997	6.4	4.7	36.2	3.9	7.0	-44.3	2.3	1.6	43.8
1998	8.0	4.9	63.3	4.1	6.4	-35.9	2.2	1.6	37.5
1999	7.3	5.4	35.2	3.3	5.9	-44.1	2.2	1.9	15.8
2000	7.7	5.7	35.1	2.7	5.8	-53.4	2.7	2.0	35.0
2001	8.0	5.9	35.6	3.1	6.0	-48.3	2.6	1.9	35.8
2002	8.3	5.9	40.7	3.1	6.0	-48.3	3.0	2.0	47.5
2003	8.4	6.2	35.5	2.5	6.0	-58.3	3.1	2.0	55.5
2004	8.6	6.1	41.0	3.1	5.8	-46.6	2.9	1.9	54.7
2005	7.7	6.4	20.3	2.9	6.1	-52.5	2.8	1.9	45.8
2006	8.7	6.3	38.1	3.0	6.2	-51.6	2.9	1.9	52.6
2007	8.2	6.4	28.1	2.1	6.1	-65.6	2.3	2.0	15.0
2008	8.7	6.4	35.9	2.6	5.9	-55.9	3.0	NA	NA

Year	Arteriosclerosis			Viral Hepatitis			HIV/AIDS		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1985	17.5	12.2	43.4	0.4	0.3	33.3	NA	NA	NA
1990	11.3	8.2	37.8	0.8	0.5	60.0	7.2	10.6	-32.1
1995	9.0	6.6	36.4	1.5	0.9	66.7	11.5	16.8	-31.4
1996	7.5	6.4	17.2	1.1	1.0	10.0	7.6	11.9	-36.1
1997	6.9	6.0	15.0	1.4	1.1	27.3	3.2	6.2	-48.1
1998	6.5	5.6	16.1	1.6	1.3	23.1	2.3	4.9	-52.8
1999	5.6	5.5	1.8	1.3	1.8	-27.8	2.2	5.3	-58.5
2000	6.4	5.2	23.1	2.2	1.9	15.8	1.8	5.2	-65.4
2001	5.3	5.0	6.0	2.5	2.0	25.0	1.9	5.0	-62.0
2002	5.7	4.8	18.8	3.5	2.0	75.0	2.5	4.9	-49.0
2003	5.5	4.4	25.0	2.6	1.8	44.4	2.5	4.7	-46.8
2004	4.6	3.9	17.9	2.9	1.8	61.1	1.8	4.5	-60.0
2005	4.8	3.8	26.3	2.3	1.8	27.8	1.5	4.2	-64.3
2006	2.8	2.7	3.7	2.2	2.3	-4.3	1.4	4.0	-65.0
2007	3.0	2.5	20.0	4.2	2.3	82.6	1.5	3.7	-59.5
2008	2.2	2.3	-4.3	3.8	2.3	65.2	1.0	3.3	-69.7

Note: US age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database. Unlike the data shown in Table 6-55, all Oregon data are from state mortality files, except for 1980 and 1985 which are from the CDC's online database. Consequently, the rates and percentage differences shown here will vary from those in Table 6-55 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. National comparability ratios have been applied to 1980-1998 rates to control for coding changes resulting from the implementation of ICD-10 when it supplanted ICD-9 in 1999. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's Disease and Alzheimer's dementia). Some differences between Oregon and US rates (e.g., alcohol-induced deaths) result, at least in part, from the state's query program (when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information).