

**TABLE 7-10. Postneonatal Deaths¹ by Weeks of Gestation and Weight
Oregon Residents, Birth Cohort 2008**

| Birthweight (In Grams) | Total | Weeks of Gestation | | | | | | | | | |
|---------------------------|-------|--------------------|-------|-------|-------|-------|----|-------|----|-----|------|
| | | <20 | 20-23 | 24-27 | 28-31 | 32-35 | 36 | 37-39 | 40 | 41+ | N.S. |
| Total | 91 | 1 | 1 | 10 | 4 | 10 | 3 | 44 | 16 | 2 | – |
| 001-349 | – | – | – | – | – | – | – | – | – | – | – |
| 350-499 | – | – | – | – | – | – | – | – | – | – | – |
| <500 | – | – | – | – | – | – | – | – | – | – | – |
| 500-749 | 8 | – | 1 | 6 | 1 | – | – | – | – | – | – |
| 750-999 | 4 | 1 | – | 3 | – | – | – | – | – | – | – |
| 1000-1249 | 2 | – | – | 1 | 1 | – | – | – | – | – | – |
| 1250-1499 | 3 | – | – | – | 2 | 1 | – | – | – | – | – |
| 1500-1999 | 1 | – | – | – | – | 1 | – | – | – | – | – |
| 2000-2499 | 12 | – | – | – | – | 5 | 2 | 5 | – | – | – |
| <2500 | 30 | 1 | 1 | 10 | 4 | 7 | 2 | 5 | – | – | – |
| 2500+ | 61 | – | – | – | – | 3 | 1 | 39 | 16 | 2 | – |
| 2500-2999 | 17 | – | – | – | – | 3 | 1 | 11 | 2 | – | – |
| 3000-3499 | 26 | – | – | – | – | – | – | 20 | 5 | 1 | – |
| 3500-3999 | 16 | – | – | – | – | – | – | 7 | 8 | 1 | – |
| 4000-4499 | 2 | – | – | – | – | – | – | 1 | 1 | – | – |
| 4500+ | – | – | – | – | – | – | – | – | – | – | – |
| Unknown | – | – | – | – | – | – | – | – | – | – | – |

– Quantity is zero.

¹ Postneonatal deaths occur from day 28 through 364 after birth.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.