

**TABLE 7-8. Early Neonatal Deaths<sup>1</sup> by Weeks of Gestation and Weight  
Oregon Residents, Birth Cohort 2009**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
<b>Total .....</b>	131	12	44	23	12	14	5	15	4	1	1
001-349 .....	24	12	11	1	—	—	—	—	—	—	—
350-499 .....	18	—	15	3	—	—	—	—	—	—	—
<500 .....	42	12	26	4	—	—	—	—	—	—	—
500-749 .....	33	—	18	14	1	—	—	—	—	—	—
750-999 .....	4	—	—	4	—	—	—	—	—	—	—
1000-1249 .....	4	—	—	1	2	1	—	—	—	—	—
1250-1499 .....	3	—	—	—	2	1	—	—	—	—	—
1500-1999 .....	15	—	—	—	7	5	1	2	—	—	—
2000-2499 .....	9	—	—	—	—	5	2	1	1	—	—
<2500 .....	110	12	44	23	12	12	3	3	1	—	—
2500+ .....	20	—	—	—	—	2	2	12	3	1	—
2500-2999 .....	8	—	—	—	—	1	1	5	1	—	—
3000-3499 .....	6	—	—	—	—	1	—	4	1	—	—
3500-3999 .....	5	—	—	—	—	—	1	3	—	1	—
4000-4499 .....	1	—	—	—	—	—	—	—	1	—	—
4500+ .....	—	—	—	—	—	—	—	—	—	—	—
Unknown .....	1	—	—	—	—	—	—	—	—	—	1

— Quantity is zero.

<sup>1</sup> Early neonatal deaths occur through day 6 after birth.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.