

**TABLE 7-8. Early Neonatal Deaths¹ by Weeks of Gestation and Weight
Oregon Residents, Birth Cohort 2010**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total ²	121	6	50	17	10	10	8	12	1	4	3
001-349	18	6	10	–	1	–	–	–	–	–	1
350-499	22	–	22	–	–	–	–	–	–	–	–
<500	40	6	32	–	1	–	–	–	–	–	1
500-749	29	–	17	11	–	–	–	–	–	–	1
750-999	7	–	1	5	1	–	–	–	–	–	–
1000-1249	7	–	–	1	4	2	–	–	–	–	–
1250-1499	1	–	–	–	–	–	1	–	–	–	–
1500-1999	13	–	–	–	4	5	3	1	–	–	–
2000-2499	6	–	–	–	–	2	1	3	–	–	–
<2500	103	6	50	17	10	9	5	4	–	–	2
2500+	17	–	–	–	–	1	3	8	1	4	–
2500-2999	4	–	–	–	–	–	1	3	–	–	–
3000-3499	7	–	–	–	–	1	2	1	1	2	–
3500-3999	6	–	–	–	–	–	–	4	–	2	–
4000-4499	–	–	–	–	–	–	–	–	–	–	–
4500+	–	–	–	–	–	–	–	–	–	–	–

¹ Early neonatal deaths occur through day 6 after birth.

² Includes unknown weight.

– Quantity is zero.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.