

**TABLE 7-9. Late Neonatal Deaths<sup>1</sup> by Weeks of Gestation and Weight  
Oregon Residents, Birth Cohort 2010**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total <sup>2</sup> .....	34	–	1	10	–	2	2	17	2	–	–
001-349 .....	–	–	–	–	–	–	–	–	–	–	–
350-499 .....	1	–	–	1	–	–	–	–	–	–	–
<500 .....	1	–	–	1	–	–	–	–	–	–	–
500-749 .....	6	–	1	5	–	–	–	–	–	–	–
750-999 .....	4	–	–	4	–	–	–	–	–	–	–
1000-1249 .....	–	–	–	–	–	–	–	–	–	–	–
1250-1499 .....	–	–	–	–	–	–	–	–	–	–	–
1500-1999 .....	–	–	–	–	–	–	–	–	–	–	–
2000-2499 .....	5	–	–	–	–	1	1	3	–	–	–
<2500 .....	16	–	1	10	–	1	1	3	–	–	–
2500+ .....	18	–	–	–	–	1	1	14	2	–	–
2500-2999 .....	10	–	–	–	–	1	1	8	–	–	–
3000-3499 .....	3	–	–	–	–	–	–	3	–	–	–
3500-3999 .....	5	–	–	–	–	–	–	3	2	–	–
4000-4499 .....	–	–	–	–	–	–	–	–	–	–	–
4500+ .....	–	–	–	–	–	–	–	–	–	–	–

<sup>1</sup> Late neonatal deaths occur from day 7 through 27 after birth.

<sup>2</sup> Includes unknown weight.

– Quantity is zero.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.