

**TABLE 7-10. Postneonatal Deaths¹ by Weeks of Gestation and Weight
Oregon Residents, Birth Cohort 2011**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total ²	63	—	—	9	2	4	2	34	9	3	—
001-349	—	—	—	—	—	—	—	—	—	—	—
350-499	1	—	—	1	—	—	—	—	—	—	—
<500	1	—	—	1	—	—	—	—	—	—	—
500-749	6	—	—	6	—	—	—	—	—	—	—
750-999	1	—	—	—	1	—	—	—	—	—	—
1000-1249	3	—	—	2	1	—	—	—	—	—	—
1250-1499	—	—	—	—	—	—	—	—	—	—	—
1500-1999	2	—	—	—	—	1	1	—	—	—	—
2000-2499	7	—	—	—	—	2	1	4	—	—	—
<2500	20	—	—	9	2	3	2	4	—	—	—
2500+	42	—	—	—	—	1	—	29	9	3	—
2500-2999	8	—	—	—	—	1	—	5	1	1	—
3000-3499	18	—	—	—	—	—	—	13	3	2	—
3500-3999	12	—	—	—	—	—	—	9	3	—	—
4000-4499	4	—	—	—	—	—	—	2	2	—	—
4500+	—	—	—	—	—	—	—	—	—	—	—

¹ Postneonatal deaths occur from day 28 through 364 after birth.

² Includes unknown weight.

— Quantity is zero.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.