

TABLE 7-8. Early neonatal deaths¹ by weeks of gestation and weight, Oregon residents, birth cohort 2013

Birthweight (in grams)	Total	Weeks of gestation								
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+
Total ²	128	12	52	32	3	9	3	10	3	4
001-349	20	11	9	—	—	—	—	—	—	—
350-499	28	—	23	4	1	—	—	—	—	—
<500	48	11	32	4	1	—	—	—	—	—
500-749	38	—	20	18	—	—	—	—	—	—
750-999	9	—	—	8	—	1	—	—	—	—
1000-1249	2	—	—	1	1	—	—	—	—	—
1250-1499	1	—	—	—	1	—	—	—	—	—
1500-1999	1	—	—	—	—	1	—	—	—	—
2000-2499	5	—	—	—	—	4	1	—	—	—
<2500	104	11	52	31	3	6	1	—	—	—
2500+	22	—	—	—	—	3	2	10	3	4
2500-2999	7	—	—	—	—	3	1	3	—	—
3000-3499	5	—	—	—	—	—	1	2	1	1
3500-3999	6	—	—	—	—	—	—	3	2	1
4000-4499	3	—	—	—	—	—	—	1	—	2
4500+	1	—	—	—	—	—	—	1	—	—

¹ Early neonatal deaths occur through day six after birth.

² Includes unknown weight.

— Quantity is zero.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.