

TABLE 7-9. Late neonatal deaths¹ by weeks of gestation and weight, Oregon residents, birth cohort 2013

Birthweight (in grams)	Total	Weeks of gestation								
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+
Total ²	35	—	—	12	5	—	1	12	3	2
001-349	—	—	—	—	—	—	—	—	—	—
350-499	2	—	—	2	—	—	—	—	—	—
<500	2	—	—	2	—	—	—	—	—	—
500-749	8	—	—	7	1	—	—	—	—	—
750-999	3	—	—	2	1	—	—	—	—	—
1000-1249	2	—	—	—	2	—	—	—	—	—
1250-1499	1	—	—	1	—	—	—	—	—	—
1500-1999	—	—	—	—	—	—	—	—	—	—
2000-2499	3	—	—	—	1	—	—	2	—	—
<2500	19	—	—	12	5	—	—	2	—	—
2500+	16	—	—	—	—	—	1	10	3	2
2500-2999	5	—	—	—	—	—	1	3	1	—
3000-3499	6	—	—	—	—	—	—	4	1	1
3500-3999	5	—	—	—	—	—	—	3	1	1
4000-4499	—	—	—	—	—	—	—	—	—	—
4500+	—	—	—	—	—	—	—	—	—	—

¹ Late neonatal deaths occur from day seven through 27 after birth.

² Includes unknown weight.

— Quantity is zero.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.