

TABLE 7-10. Postneonatal deaths¹ by weeks of gestation and weight, Oregon residents, birth cohort 2013

Birthweight (in grams)	Total	Weeks of gestation								
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+
Total ²	69	–	–	7	6	8	5	33	8	2
001-349	–	–	–	–	–	–	–	–	–	–
350-499	–	–	–	–	–	–	–	–	–	–
<500	–	–	–	–	–	–	–	–	–	–
500-749	5	–	–	2	1	2	–	–	–	–
750-999	1	–	–	1	–	–	–	–	–	–
1000-1249	5	–	–	1	3	1	–	–	–	–
1250-1499	4	–	–	3	1	–	–	–	–	–
1500-1999	4	–	–	–	1	1	1	1	–	–
2000-2499	5	–	–	–	–	3	–	2	–	–
<2500	24	–	–	7	6	7	1	3	–	–
2500+	45	–	–	–	–	1	4	30	8	2
2500-2999	19	–	–	–	–	1	4	11	2	1
3000-3499	16	–	–	–	–	–	–	12	4	–
3500-3999	7	–	–	–	–	–	–	5	1	1
4000-4499	2	–	–	–	–	–	–	1	1	–
4500+	1	–	–	–	–	–	–	1	–	–

¹ Postneonatal deaths occur from day 28 through 364 after birth.

² Includes unknown weight.

– Quantity is zero.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.