

TABLE 6-58. Age-adjusted death rates for selected causes of death, Oregon and United States residents, 2000-2014¹

| Year | Total | | | Cancer | | | Heart disease | | |
|------------|--------|-------|--------|--------|-------|--------|---------------|-------|--------|
| | Oregon | US | % Diff | Oregon | US | % Diff | Oregon | US | % Diff |
| 2000 | 826.9 | 869.0 | -4.8 | 197.6 | 199.6 | -1.0 | 197.5 | 257.6 | -23.3 |
| 2001 | 835.9 | 851.6 | -1.8 | 198.7 | 195.6 | 1.6 | 195.2 | 246.8 | -20.9 |
| 2002 | 855.0 | 845.3 | 1.1 | 200.9 | 193.5 | 3.8 | 198.0 | 240.8 | -17.8 |
| 2003 | 838.4 | 832.7 | 0.7 | 198.3 | 190.1 | 4.3 | 189.5 | 232.3 | -18.4 |
| 2004 | 814.8 | 800.8 | 1.7 | 196.7 | 185.8 | 5.9 | 179.2 | 217.0 | -17.4 |
| 2005 | 791.4 | 798.8 | -0.9 | 189.4 | 183.8 | 3.0 | 169.5 | 211.1 | -19.7 |
| 2006 | 784.5 | 776.5 | 1.0 | 185.7 | 180.7 | 2.8 | 162.6 | 200.2 | -18.8 |
| 2007 | 771.6 | 760.2 | 1.5 | 184.7 | 178.4 | 3.5 | 159.7 | 190.9 | -16.3 |
| 2008 | 772.8 | 758.3 | 1.9 | 182.8 | 175.3 | 4.3 | 154.5 | 186.5 | -17.2 |
| 2009 | 739.7 | 741.1 | -0.2 | 176.7 | 173.2 | 2.0 | 143.0 | 180.1 | -20.6 |
| 2010 | 735.0 | 747.0 | -1.6 | 177.9 | 172.8 | 2.9 | 139.7 | 179.1 | -22.0 |
| 2011 | 730.0 | 741.3 | -1.5 | 172.7 | 169.0 | 2.2 | 136.2 | 173.7 | -21.6 |
| 2012 | 706.4 | 732.8 | -3.6 | 167.5 | 166.5 | 0.6 | 130.3 | 170.5 | -23.6 |
| 2013 | 716.8 | 731.9 | -2.1 | 163.0 | 163.2 | -0.1 | 134.6 | 169.8 | -20.7 |
| 2014 | 702.8 | 724.6 | -3.0 | 159.3 | 161.2 | -1.2 | 131.3 | 167.0 | -21.4 |

| Year | Chronic lower resp. disease | | | Unintentional injuries | | | Cerebrovascular disease | | |
|------------|-----------------------------|------|--------|------------------------|------|--------|-------------------------|------|--------|
| | Oregon | US | % Diff | Oregon | US | % Diff | Oregon | US | % Diff |
| 2000 | 47.8 | 44.2 | 8.1 | 34.6 | 34.9 | -0.9 | 70.8 | 60.9 | 16.3 |
| 2001 | 48.7 | 43.6 | 11.7 | 35.4 | 35.5 | -0.3 | 71.4 | 57.7 | 23.7 |
| 2002 | 50.9 | 43.5 | 17.0 | 38.4 | 36.9 | 4.1 | 71.7 | 56.2 | 27.6 |
| 2003 | 49.8 | 43.3 | 15.0 | 38.3 | 37.3 | 2.7 | 68.5 | 53.5 | 28.0 |
| 2004 | 48.1 | 41.1 | 17.0 | 38.8 | 37.7 | 2.9 | 61.9 | 50.0 | 23.8 |
| 2005 | 47.8 | 43.2 | 10.6 | 37.6 | 39.1 | -3.8 | 57.3 | 46.6 | 23.0 |
| 2006 | 46.8 | 40.5 | 15.6 | 40.7 | 39.8 | 2.3 | 48.8 | 43.6 | 11.9 |
| 2007 | 47.5 | 40.8 | 16.4 | 41.7 | 40.0 | 4.3 | 44.5 | 42.2 | 5.5 |
| 2008 | 48.2 | 44.0 | 9.5 | 42.4 | 38.8 | 9.3 | 45.6 | 40.7 | 12.0 |
| 2009 | 46.4 | 42.3 | 9.6 | 38.8 | 37.3 | 3.9 | 44.0 | 38.9 | 13.2 |
| 2010 | 46.5 | 42.2 | 10.2 | 37.8 | 38.0 | -0.6 | 40.5 | 39.1 | 3.6 |
| 2011 | 45.6 | 42.5 | 7.2 | 40.4 | 39.1 | 3.4 | 42.0 | 37.9 | 10.8 |
| 2012 | 42.0 | 41.5 | 1.1 | 38.9 | 39.1 | -0.6 | 37.5 | 36.9 | 1.5 |
| 2013 | 42.6 | 42.1 | 1.2 | 39.6 | 39.4 | 0.5 | 37.0 | 36.2 | 2.3 |
| 2014 | 39.7 | 40.5 | -1.9 | 40.7 | 40.5 | 0.4 | 37.0 | 36.5 | 1.5 |

¹ Most recent year for which final US data are available.

NOTE: US age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database. Unlike the data shown in tables 6-54 and 6-55, all Oregon data are from state mortality files. Consequently, the rates and percentage differences shown here will vary from those in tables 6-54 and 6-55 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's disease vs. Alzheimer's dementia). Some differences between Oregon and US rates (e.g., alcohol-induced deaths) result, at least in part, from the Oregon's query program: when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information.

TABLE 6-58. Age-adjusted death rates for selected causes of death, Oregon and United States residents, 2000-2014¹ — Continued

| Year | Alzheimer's disease | | | Diabetes mellitus | | | Suicide | | |
|------------|---------------------|------|--------|-------------------|------|--------|---------|------|--------|
| | Oregon | US | % Diff | Oregon | US | % Diff | Oregon | US | % Diff |
| 2000 | 24.8 | 18.1 | 37.0 | 23.8 | 25.0 | -4.8 | 14.3 | 10.4 | 37.5 |
| 2001 | 28.1 | 19.0 | 47.9 | 28.8 | 25.2 | 14.3 | 14.9 | 10.7 | 39.3 |
| 2002 | 30.3 | 20.2 | 50.0 | 28.6 | 25.4 | 12.6 | 14.5 | 10.9 | 33.0 |
| 2003 | 30.6 | 21.4 | 43.0 | 28.1 | 25.3 | 11.1 | 16.3 | 10.8 | 50.9 |
| 2004 | 33.4 | 21.8 | 53.2 | 29.0 | 24.5 | 18.4 | 15.2 | 10.9 | 39.4 |
| 2005 | 30.4 | 22.9 | 32.8 | 29.3 | 24.6 | 19.1 | 14.9 | 10.9 | 36.7 |
| 2006 | 29.5 | 22.6 | 30.5 | 28.9 | 23.3 | 24.0 | 15.1 | 10.9 | 38.5 |
| 2007 | 28.0 | 22.7 | 23.3 | 27.9 | 22.5 | 24.0 | 15.6 | 11.3 | 38.1 |
| 2008 | 30.5 | 24.4 | 25.0 | 24.8 | 21.8 | 13.8 | 14.7 | 11.6 | 26.7 |
| 2009 | 27.7 | 23.5 | 17.8 | 25.3 | 20.9 | 20.9 | 16.1 | 11.8 | 36.2 |
| 2010 | 28.7 | 25.1 | 14.3 | 24.2 | 20.8 | 16.3 | 17.1 | 12.1 | 41.4 |
| 2011 | 28.8 | 24.7 | 16.7 | 24.8 | 21.6 | 14.9 | 16.2 | 12.3 | 31.4 |
| 2012 | 28.1 | 23.8 | 18.2 | 24.4 | 21.2 | 14.9 | 17.6 | 12.6 | 40.0 |
| 2013 | 27.1 | 23.5 | 15.4 | 23.4 | 21.2 | 10.6 | 16.8 | 12.6 | 33.4 |
| 2014 | 28.3 | 25.4 | 11.3 | 22.3 | 20.9 | 6.5 | 18.6 | 13.0 | 43.1 |

| Year | Alcohol-induced | | | Hypertension | | | Flu & pneumonia | | |
|------------|-----------------|-----|--------|--------------|-----|--------|-----------------|------|--------|
| | Oregon | US | % Diff | Oregon | US | % Diff | Oregon | US | % Diff |
| 2000 | 10.8 | 7.0 | 54.3 | 6.2 | 6.5 | -4.6 | 17.5 | 23.7 | -26.2 |
| 2001 | 12.2 | 7.0 | 74.3 | 8.6 | 6.8 | 26.5 | 15.7 | 21.9 | -28.3 |
| 2002 | 12.3 | 6.9 | 78.3 | 9.6 | 7.0 | 37.1 | 17.9 | 22.6 | -20.8 |
| 2003 | 14.2 | 7.0 | 102.9 | 9.3 | 7.4 | 25.7 | 17.0 | 22.0 | -22.7 |
| 2004 | 13.8 | 7.0 | 97.1 | 9.5 | 7.7 | 23.4 | 14.7 | 19.8 | -25.8 |
| 2005 | 13.7 | 7.0 | 95.7 | 10.6 | 8.0 | 32.5 | 15.1 | 20.3 | -25.6 |
| 2006 | 11.7 | 7.0 | 67.1 | 8.9 | 7.5 | 18.7 | 12.8 | 17.8 | -28.1 |
| 2007 | 13.1 | 7.3 | 79.5 | 8.6 | 7.4 | 16.2 | 11.4 | 16.2 | -29.6 |
| 2008 | 12.9 | 7.4 | 74.3 | 9.5 | 7.7 | 23.4 | 12.3 | 16.9 | -27.2 |
| 2009 | 13.4 | 7.4 | 81.4 | 9.5 | 7.7 | 23.1 | 12.0 | 16.2 | -26.2 |
| 2010 | 13.0 | 7.6 | 71.2 | 9.8 | 8.0 | 23.1 | 9.3 | 15.1 | -38.4 |
| 2011 | 14.6 | 7.7 | 89.3 | 9.7 | 8.1 | 19.5 | 8.7 | 15.7 | -44.5 |
| 2012 | 14.7 | 8.0 | 84.3 | 10.4 | 8.2 | 27.2 | 8.1 | 14.4 | -43.8 |
| 2013 | 15.4 | 8.2 | 88.2 | 10.7 | 8.5 | 25.9 | 10.5 | 15.9 | -34.1 |
| 2014 | 16.4 | 8.5 | 93.2 | 9.8 | 8.2 | 20.0 | 9.1 | 15.1 | -39.9 |

¹ Most recent year for which final US data are available.

NOTE: US age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database. Unlike the data shown in tables 6-54 and 6-55, all Oregon data are from state mortality files. Consequently, the rates and percentage differences shown here will vary from those in tables 6-54 and 6-55 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's disease vs. Alzheimer's dementia). Some differences between Oregon and US rates (e.g., alcohol-induced deaths) result, at least in part, from the Oregon's query program: when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information.

TABLE 6-58. Age-adjusted death rates for selected causes of death, Oregon and United States residents, 2000-2014¹ — Continued

| Year | Parkinson's disease | | | Viral hepatitis | | | Amyotrophic lateral sclerosis | | |
|------------|---------------------|-----|--------|-----------------|-----|--------|-------------------------------|-----|--------|
| | Oregon | US | % Diff | Oregon | US | % Diff | Oregon | US | % Diff |
| 2000 | 7.7 | 5.7 | 35.1 | 2.2 | 1.9 | 15.8 | 2.7 | 2.0 | 35.0 |
| 2001 | 8.0 | 5.8 | 37.9 | 2.5 | 2.0 | 25.0 | 2.6 | 1.9 | 36.8 |
| 2002 | 8.3 | 5.9 | 40.7 | 3.5 | 2.0 | 75.0 | 3.0 | 2.0 | 50.0 |
| 2003 | 8.4 | 6.2 | 35.5 | 2.6 | 1.8 | 44.4 | 3.1 | 2.0 | 55.0 |
| 2004 | 8.6 | 6.1 | 41.0 | 2.9 | 1.8 | 61.1 | 2.9 | 1.9 | 52.6 |
| 2005 | 7.7 | 6.4 | 20.3 | 2.3 | 1.8 | 27.8 | 2.8 | 1.9 | 47.4 |
| 2006 | 8.7 | 6.3 | 38.1 | 2.2 | 2.3 | -4.3 | 2.9 | 1.9 | 52.6 |
| 2007 | 8.2 | 6.4 | 28.1 | 4.2 | 2.3 | 82.6 | 2.3 | 1.9 | 21.1 |
| 2008 | 8.7 | 6.4 | 35.9 | 3.8 | 2.3 | 65.2 | 3.0 | 1.9 | 57.9 |
| 2009 | 8.3 | 6.4 | 29.7 | 3.9 | 2.2 | 77.1 | 2.7 | 1.9 | 39.8 |
| 2010 | 8.5 | 6.8 | 25.6 | 3.8 | 2.1 | 79.8 | 2.8 | 2.0 | 38.9 |
| 2011 | 8.0 | 7.0 | 14.1 | 3.8 | 2.2 | 74.7 | 2.7 | 2.0 | 33.8 |
| 2012 | 8.0 | 7.0 | 14.3 | 3.2 | 2.1 | 51.3 | 2.7 | 2.1 | 29.4 |
| 2013 | 8.5 | 7.3 | 15.9 | 4.6 | 2.1 | 118.1 | 2.9 | 2.0 | 45.6 |
| 2014 | 8.0 | 7.4 | 7.7 | 4.1 | 2.1 | 95.8 | 2.6 | 2.0 | 30.7 |

| Year | Homicide | | | HIV/AIDS | | | Arteriosclerosis | | |
|------------|----------|-----|--------|----------|-----|--------|------------------|-----|--------|
| | Oregon | US | % Diff | Oregon | US | % Diff | Oregon | US | % Diff |
| 2000 | 2.7 | 5.9 | -54.2 | 1.8 | 5.2 | -65.4 | 6.4 | 5.2 | 23.1 |
| 2001 | 3.1 | 7.1 | -56.3 | 1.9 | 5.0 | -62.0 | 5.3 | 5.0 | 6.0 |
| 2002 | 3.1 | 6.1 | -49.2 | 2.5 | 4.9 | -49.0 | 5.7 | 4.7 | 21.3 |
| 2003 | 2.5 | 6.0 | -58.3 | 2.5 | 4.7 | -46.8 | 5.5 | 4.4 | 25.0 |
| 2004 | 3.1 | 5.9 | -47.5 | 1.8 | 4.5 | -60.0 | 4.6 | 3.9 | 17.9 |
| 2005 | 2.9 | 6.1 | -52.5 | 1.5 | 4.2 | -64.3 | 4.8 | 3.8 | 26.3 |
| 2006 | 3.0 | 6.2 | -51.6 | 1.4 | 4.0 | -65.0 | 2.8 | 2.7 | 3.7 |
| 2007 | 2.1 | 6.1 | -65.6 | 1.5 | 3.7 | -59.5 | 3.0 | 2.5 | 20.0 |
| 2008 | 2.6 | 5.9 | -55.9 | 1.0 | 3.3 | -69.7 | 2.2 | 2.3 | -4.3 |
| 2009 | 2.6 | 5.5 | -53.3 | 1.1 | 3.0 | -62.7 | 1.8 | 2.2 | -19.0 |
| 2010 | 2.9 | 5.3 | -45.5 | 1.2 | 2.6 | -55.1 | 1.6 | 2.2 | -29.1 |
| 2011 | 2.8 | 5.3 | -47.5 | 0.9 | 2.4 | -62.7 | 2.0 | 2.0 | -0.5 |
| 2012 | 2.8 | 5.4 | -47.9 | 1.4 | 2.2 | -37.8 | 1.1 | 2.0 | -44.9 |
| 2013 | 2.3 | 5.2 | -56.3 | 1.2 | 2.1 | -43.5 | 1.2 | 1.8 | -33.8 |
| 2014 | 2.4 | 5.1 | -52.3 | 0.8 | 2.0 | -60.9 | 0.8 | 1.7 | -51.8 |

¹ Most recent year for which final US data are available.

NOTE: US age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database. Unlike the data shown in tables 6-54 and 6-55, all Oregon data are from state mortality files. Consequently, the rates and percentage differences shown here will vary from those in tables 6-54 and 6-55 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's disease vs. Alzheimer's dementia). Some differences between Oregon and US rates (e.g., alcohol-induced deaths) result, at least in part, from the Oregon's query program: when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information.