

TABLE 7-10. Postneonatal deaths¹ by weeks of gestation and weight, Oregon residents, birth cohort 2014

Birthweight (in grams)	Total	Weeks of gestation ²								
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+
Total ³	73	–	–	4	3	9	4	34	13	6
001-349	–	–	–	–	–	–	–	–	–	–
350-499	1	–	–	1	–	–	–	–	–	–
<500	1	–	–	1	–	–	–	–	–	–
500-749	1	–	–	1	–	–	–	–	–	–
750-999	1	–	–	1	–	–	–	–	–	–
1000-1249	2	–	–	1	1	–	–	–	–	–
1250-1499	2	–	–	–	1	1	–	–	–	–
1500-1999	3	–	–	–	1	2	–	–	–	–
2000-2499	7	–	–	–	–	4	2	–	–	1
<2500	17	–	–	4	3	7	2	–	–	1
2500+	55	–	–	–	–	2	2	33	13	5
2500-2999	16	–	–	–	–	2	1	10	2	1
3000-3499	18	–	–	–	–	–	1	12	5	–
3500-3999	17	–	–	–	–	–	–	9	4	4
4000-4499	4	–	–	–	–	–	–	2	2	–
4500+	–	–	–	–	–	–	–	–	–	–

– Quantity is zero.

¹ Postneonatal deaths occur from day 28 through 364 after birth.

² Clinical estimate of gestation. If clinical estimate is unknown, gestation is calculated from reported date of last menses.

³ Total includes reports with unknown birthweight.