

TABLE 6-35. Fatal overdoses and poisonings by manner, type, sex, and age, Oregon residents, 2017

Manner and type of substance ¹	Total*	M	F	Age at death									
				0-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+
Total	1,052	694	357	2	1	58	136	148	230	264	154	38	21
Mental and behavioral disorders due to psychoactive substance use²	434	309	125	–	–	4	22	30	103	133	106	23	13
Alcohol ³	291	210	81	–	–	–	9	20	69	98	75	15	5
Opioids	22	11	11	–	–	–	5	2	4	5	2	1	3
Cannabinoids	–	–	–	–	–	–	–	–	–	–	–	–	–
Sedatives and hypnotics	–	–	–	–	–	–	–	–	–	–	–	–	–
Cocaine	3	2	1	–	–	–	–	–	1	1	1	–	–
Other stimulants	36	26	10	–	–	–	1	4	15	11	5	–	–
Hallucinogens	–	–	–	–	–	–	–	–	–	–	–	–	–
Tobacco ⁴	54	44	10	–	–	–	–	1	8	15	20	6	4
Volatile solvents	–	–	–	–	–	–	–	–	–	–	–	–	–
Other (multiple) psychoactive substances	28	16	12	–	–	4	7	3	6	3	3	1	1
Unintentional overdoses/poisoning	450	302	148	–	–	42	94	92	92	93	27	4	6
Nonopioid analgesics, antipyretics, etc.	5	2	3	–	–	–	–	2	1	–	–	1	1
Psychotropic, sedative-hypnotic drugs	84	62	22	–	–	5	12	14	21	26	5	1	–
Narcotics and hallucinogens ⁵	167	115	52	–	–	25	57	27	25	24	9	–	–
Other and unspecified drugs ⁶	145	87	58	–	–	10	19	41	33	29	8	1	4
Alcohol	35	25	10	–	–	1	4	4	11	10	5	–	–
Organic solvents & halogenated HC ⁷	1	1	–	–	–	–	–	1	–	–	–	–	–
Carbon monoxide & other gases	11	9	2	–	–	1	1	3	1	3	–	1	1
Pesticides	1	–	1	–	–	–	–	–	–	1	–	–	–
Other chemicals & substances	1	1	–	–	–	–	1	–	–	–	–	–	–
<i>Any opioid</i> ⁸	274	180	94	–	–	32	69	57	49	53	13	–	1
Intentional self-poisoning	130	67	62	–	1	8	18	22	28	26	16	9	2
Nonopioid analgesics, antipyretics, etc.	1	1	–	–	–	–	–	1	–	–	–	–	–
Psychotropic, sedative-hypnotic drugs	17	6	11	–	–	–	6	1	4	4	2	–	–
Narcotics and hallucinogens ⁵	18	6	11	–	1	1	2	–	3	6	3	2	–
Other and unspecified drugs ⁶	55	25	30	–	–	4	8	13	11	8	5	6	–
Alcohol	3	2	1	–	–	–	–	1	–	2	–	–	–
Organic solvents & halogenated HC ⁷	5	1	4	–	–	–	–	2	1	1	1	–	–
Carbon monoxide & other gases	30	25	5	–	–	3	2	4	9	5	5	1	1
Pesticides	–	–	–	–	–	–	–	–	–	–	–	–	–
Other chemicals & substances	1	1	–	–	–	–	–	–	–	–	–	–	1
<i>Any opioid</i> ⁸	41	16	24	–	1	2	6	4	5	10	7	6	–

See footnotes at end of table.

TABLE 6-35. Fatal overdoses and poisonings by manner, type, sex, and age, Oregon residents, 2017 — Continued

Manner and type of substance ¹	Total*	M	F	Age at death										
				0-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	
Assault by poisoning	1	—	1	1	—	—	—	—	—	—	—	—	—	—
Undetermined intent	37	16	21	1	—	4	2	4	7	12	5	2	—	—
Nonopioid analgesics, antipyretics, etc.	1	—	1	—	—	—	—	—	1	—	—	—	—	—
Psychotropic, sedative-hypnotic drugs	11	5	6	—	—	2	1	1	1	4	2	—	—	—
Narcotics and hallucinogens ⁵	11	5	6	1	—	1	1	1	1	4	1	1	—	—
Other and unspecified drugs ⁶	9	3	6	—	—	—	—	2	3	3	1	—	—	—
Alcohol	2	1	1	—	—	—	—	—	1	1	—	—	—	—
Organic solvents & halogenated HC ⁷	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Carbon monoxide & other gases	2	2	—	—	—	—	—	—	—	—	1	1	—	—
Pesticides	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other chemicals & substances	1	—	1	—	—	1	—	—	—	—	—	—	—	—
<i>Any opioid</i> ⁸	18	7	11	1	—	1	1	3	4	5	2	1	—	—

— Quantity is zero.

* Includes unknown sex.

¹ The distinction between deaths classified as mental/behavioral disorders due to psychoactive substance use versus injury deaths is somewhat factitious. Deaths attributed to drug toxicity are classified to the former category while deaths attributed to poisoning are classified as injury deaths. If the certifying physician notes that a death is due to chronic drug use, then the death is classified to mental/behavioral disorders, but this may not be done in all applicable cases. Other "natural" causes, such as drug-induced hypopituitarism, are not included here but are included in tables 6-6, 6-7, 6-17, and 6-18, among others.

² Includes ICD-10 codes F10-F19.

³ Most deaths involving abusive alcohol use are attributed to other organ systems (e.g., alcoholic cirrhosis of the liver). See "Alcohol-induced deaths" in other tables, such as 6-6, 6-7, 6-17, and 6-18, for a more inclusive count. Note these figures, too, are undercounts, as they do not include injury deaths in which alcohol played a critical role (e.g., motor vehicle crashes, homicides).

⁴ Most deaths resulting from tobacco use were attributed to other organ systems (e.g., lung cancer, emphysema, heart disease). See tables 6-19 through 6-21 for a more complete account of tobacco-linked deaths.

⁵ Includes other drugs acting on the autonomic nervous system.

⁶ Includes deaths due to poisoning from multiple substances in more than one category.

⁷ HC = hydrocarbons.

⁸ Fatal overdoses/poisonings in which a specific opioid was coded on the death certificate (ICD-10 codes T40.0-T40.4, T40.6). Some death certificates involving overdoses/poisonings do not mention the specific drug(s). Category is not mutually exclusive.