

TABLE 6-55. Age-adjusted death rates for residents of Oregon and the United States for leading causes of death, 2016¹

| Cause | Age-adjusted rate ² | | Percent difference | State rank ³ | ICD-10 codes ⁴ |
|---|--------------------------------|--------|--------------------|-------------------------|---|
| | U.S. | Oregon | | | |
| All causes | 728.8 | 705.9 | -3.1 | 38 | A00-Y89.9 |
| Malignant neoplasms | 155.8 | 155.9 | 0.1 | 32 | C00-C97 |
| Heart disease | 165.5 | 135.0 | -18.4 | 47 | I00-I09, I11, I13, I20-I51 |
| Unintended injuries | 47.4 | 46.0 | -3.0 | 34 | V01-X59, Y85-Y86 |
| Chronic lower respiratory disease | 40.6 | 40.4 | -0.5 | 30 | J40-J47 |
| Cerebrovascular disease | 37.3 | 37.8 | 1.3 | 25 | I60-I69 |
| Alzheimer's disease | 30.3 | 34.8 | 14.9 | 20 | G30 |
| Diabetes mellitus | 21.0 | 24.0 | 14.3 | 11 | E10-E14 |
| Suicide | 13.5 | 17.8 | 31.9 | 16 | X60-X84, Y87.0 |
| Alcohol-induced deaths | 9.5 | 16.9 | 77.9 | 5 | E24.4, F10, G31.2, G62.1, G72.1, I42.6, K29.2, K70, K86.0, R78.0, X45, X65, Y15 |
| Hypertension | 8.6 | 10.6 | 23.3 | 8 | I10, I12, I15 |
| Parkinson's disease | 8.0 | 9.1 | 13.8 | 7 | G20-G21 |
| Influenza & pneumonia | 13.5 | 8.9 | -34.1 | 49 | J09-J18 |
| Nephritis & nephrosis | 13.1 | 7.7 | -41.2 | 47 | N00-N07, N17-N19, N25-N27 |
| Septicemia | 10.7 | 5.2 | -51.4 | 49 | A40-A41 |
| Congenital anomalies | 3.2 | 3.3 | 3.1 | 30 | Q00-Q99 |
| Homicide | 6.2 | 3.3 | -46.8 | 37 | X85-Y09, Y87.1 |
| Perinatal conditions | 4.0 | 3.0 | -25.0 | 44 | P00-P96 |
| Viral hepatitis | 1.6 | 2.9 | 81.3 | 4 | B15-B19 |
| Aortic aneurysm & dissection | 2.6 | 2.5 | -3.8 | 34 | I71 |
| Amyotrophic lateral sclerosis | 1.7 | 2.1 | 23.5 | 14 | G12.2 |
| Arteriosclerosis | 1.5 | 0.9 | -40.0 | 32 | I70 |
| HIV/AIDS | 1.8 | 0.8 | -55.6 | 29 | B20-B24 |

¹ Most recent year for which final data are available.

² Rates are adjusted to the U.S. standard million population and are per 100,000. Age-adjusted death rates allow the comparison of Oregon and the U.S. as if the population structure of each were identical (Oregon's population is older than the U.S. as a whole). All rates in this table were calculated using the federal Centers for Disease Control and Prevention WONDER (Wide-ranging Online Data for Epidemiological Research) system (<http://wonder.cdc.gov>). These rates may vary slightly from rates published by the National Center for Health Statistics and the Oregon Center for Health Statistics due to different file closure dates and different population estimate methodologies.

³ Ranked from high (1) to low (51) among the 50 states and the District of Columbia. Rankings for some causes of death are not out of a total of 51 because states with unreliable data have been excluded.

⁴ From the World Health Organization's International Classification of Disease, 10th Edition.