

TABLE 7-10. Postneonatal deaths¹ by weeks of gestation and weight, Oregon residents, birth cohort 2016

Birthweight (in grams)	Total	Weeks of gestation ²								
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+
Total	65	–	3	3	2	9	2	30	8	8
001-349	–	–	–	–	–	–	–	–	–	–
350-499	–	–	–	–	–	–	–	–	–	–
<500	–	–	–	–	–	–	–	–	–	–
500-749	6	–	3	2	1	–	–	–	–	–
750-999	1	–	–	1	–	–	–	–	–	–
1000-1249	–	–	–	–	–	–	–	–	–	–
1250-1499	–	–	–	–	–	–	–	–	–	–
1500-1999	7	–	–	–	1	5	1	–	–	–
2000-2499	7	–	–	–	–	1	–	6	–	–
<2500	21	–	3	3	2	6	1	6	–	–
2500+	44	–	–	–	–	3	1	24	8	8
2500-2999	12	–	–	–	–	3	–	5	2	2
3000-3499	25	–	–	–	–	–	1	15	5	4
3500-3999	3	–	–	–	–	–	–	2	–	1
4000-4499	3	–	–	–	–	–	–	2	1	–
4500+	1	–	–	–	–	–	–	–	–	1

– Quantity is zero.

¹ Postneonatal deaths occur from day 28 through 364 after birth.

² Clinical estimate of gestation. If clinical estimate is unknown, gestation is calculated from reported date of last menses.