

**TABLE 13. Tobacco-linked deaths by cause of death
Oregon residents, 2018**

Selected causes of death (and their ICD-10 codes)	Total	Linked ¹		Not linked		Unknown	
		No.	%	No.	%	No.	%
Total ²	36,191	7,759	21.4	18,834	52.0	9,598	26.5
Malignant neoplasms	3,651	1,767	48.4	1,128	30.9	756	20.7
Oral cavity, lip, pharynx (C00.0-C14.8)	148	66	44.6	41	27.7	41	27.7
Esophagus (C15)	248	84	33.9	92	37.1	72	29.0
Stomach (C16)	112	23	20.5	58	51.8	31	27.7
Pancreas (C25)	620	82	13.2	384	61.9	154	24.8
Larynx (C32)	38	27	71.1	3	7.9	8	21.1
Lung, bronchi, trachea (C33-C34)	1,811	1,339	73.9	194	10.7	278	15.4
Cervix uteri (C53)	47	9	19.1	30	63.8	8	17.0
Kidney, other urinary tract (C64-C65)	214	47	22.0	115	53.7	52	24.3
Urinary bladder (C67)	261	88	33.7	90	34.5	83	31.8
Acute Myeloid Leukemia (C92.0)	152	2	1.3	121	79.6	29	19.1
Cardiovascular diseases	8,709	1,941	22.3	4,095	47.0	2,673	30.7
Ischemic heart disease (I20-I25)	3,168	976	30.8	1,265	39.9	927	29.3
Other heart disease (I00-I09, I26-I51)	3,167	547	17.3	1,676	52.9	944	29.8
Cerebrovascular disease (I60-I69)	2,027	297	14.7	1,036	51.1	694	34.2
Atherosclerosis (I70)	44	18	40.9	17	38.6	9	20.5
Aortic aneurysm (I71)	155	49	31.6	53	34.2	53	34.2
Other arterial disease (I72-I78)	148	54	36.5	48	32.4	46	31.1
Respiratory diseases	2,380	1,570	66.0	430	18.1	380	16.0
Pneumonia & influenza (J09-J18)	528	82	15.5	293	55.5	153	29.0
Bronchitis & emphysema (J40-J43)	139	112	80.6	13	9.4	14	10.1
Other chronic airways obstruction (J44)	1,713	1,376	80.3	124	7.2	213	12.4
Perinatal conditions ³	58	–	–	44	75.9	14	24.1
Selected perinatal conditions ⁴	31	–	–	26	83.9	5	16.1
Sudden infant death syndrome (R95)	27	–	–	18	66.7	9	33.3
Other causes	21,393	2,481	11.6	13,137	61.4	5,775	27.0

– Quantity is zero.

¹ The Oregon death certificate provides four possible answers to the question, "Did tobacco use contribute to death?": yes, probably, no and unknown. The linked category includes deaths listed as yes or probably, or if a contributing cause is F17.

² The causes of death shown in this table are those linked to tobacco use by the federal Centers for Disease Control and Prevention (CDC. Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses -- United

³ The number of infant deaths resulting from exposure to tobacco combustion products is underreported.

⁴ The category includes the following conditions: other disorders related to short gestation and low birthweight (P07), respiratory distress of newborn (P22), congenital pneumonia (P23), neonatal aspiration syndromes (P24), and other respiratory conditions originating in the perinatal period (P25-P28).