

**TABLE 6. Tobacco-linked deaths by cause of death
Oregon residents, 2019**

Selected causes of death (and their ICD-10 codes)	Total	Linked ¹		Not linked		Unknown	
		No.	%	No.	%	No.	%
Total ²	37,397	8,003	21.4	19,246	51.5	10,148	27.1
Malignant neoplasms	3,597	1,719	47.8	1,148	31.9	730	20.3
Oral cavity, lip, pharynx (C00.0-C14.8)	170	70	41.2	54	31.8	46	27.1
Esophagus (C15)	237	92	38.8	75	31.6	70	29.5
Stomach (C16)	126	15	11.9	91	72.2	20	15.9
Pancreas (C25)	608	73	12.0	388	63.8	147	24.2
Larynx (C32)	31	19	61.3	5	16.1	7	22.6
Lung, bronchi, trachea (C33-C34)	1,782	1,311	73.6	190	10.7	281	15.8
Cervix uteri (C53)	39	4	10.3	25	64.1	10	25.6
Kidney, other urinary tract (C64-C65)	208	42	20.2	118	56.7	48	23.1
Urinary bladder (C67)	248	82	33.1	92	37.1	74	29.8
Acute Myeloid Leukemia (C92.0)	148	11	7.4	110	74.3	27	18.2
Cardiovascular diseases	9,095	1,976	21.7	4,150	45.6	2,969	32.6
Ischemic heart disease (I20-I25)	3,390	1,007	29.7	1,289	38.0	1,094	32.3
Other heart disease (I00-I09, I26-I51)	3,230	542	16.8	1,683	52.1	1,005	31.1
Cerebrovascular disease (I60-I69)	2,134	307	14.4	1,060	49.7	767	35.9
Atherosclerosis (I70)	37	12	32.4	11	29.7	14	37.8
Aortic aneurysm (I71)	163	57	35.0	49	30.1	57	35.0
Other arterial disease (I72-I78)	141	51	36.2	58	41.1	32	22.7
Respiratory diseases	2,541	1,740	68.5	401	15.8	400	15.7
Pneumonia & influenza (J09-J18)	486	81	16.7	260	53.5	145	29.8
Bronchitis & emphysema (J40-J43)	159	138	86.8	11	6.9	10	6.3
Other chronic airways obstruction (J44)	1,896	1,521	80.2	130	6.9	245	12.9
Perinatal conditions ³	58	–	–	47	81.0	11	19.0
Selected perinatal conditions ⁴	35	–	–	33	94.3	2	5.7
Sudden infant death syndrome (R95)	23	–	–	14	60.9	9	39.1
Other causes	22,106	2,568	11.6	13,500	61.1	6,038	27.3

– Quantity is zero.

¹ The Oregon death certificate provides four possible answers to the question, "Did tobacco use contribute to death?": yes, probably, no and unknown. The linked category includes deaths listed as yes or probably, or if a contributing cause is F17.

² The causes of death shown in this table are those linked to tobacco use by the federal Centers for Disease Control and Prevention (CDC. Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses -- United States, 1997-2001.

³ The number of infant deaths resulting from exposure to tobacco combustion products is underreported.

⁴ The category includes the following conditions: other disorders related to short gestation and low birthweight (P07), respiratory distress of newborn (P22), congenital pneumonia (P23), neonatal aspiration syndromes (P24), and other respiratory conditions originating in the perinatal period (P25-P28).