

**Measures of infant birthweight, Oregon
residents, 1980-2016**

Year	Very low birthweight ¹	Low birthweight ²	Macrosomia ³
1980	8.2	42.1	13.8
1981	8.3	40.1	13.9
1982	8.5	40.6	14.1
1983	8.5	41.5	14.4
1984	8.7	42.8	14.7
1985	8.8	42.4	14.4
1986	8.3	43.0	14.7
1987	8.5	45.5	14.2
1988	7.9	44.7	14.1
1989	8.9	43.3	14.6
1990	8.2	41.9	14.2
1991	8.3	41.0	13.9
1992	8.6	43.2	13.8
1993	9.4	43.1	13.8
1994	8.6	44.4	13.8
1995	8.8	46.1	13.5
1996	9.2	44.3	13.1
1997	8.9	46.1	12.8
1998	9.0	44.7	13.0
1999	9.4	44.6	12.8
2000	9.9	46.8	12.8
2001	9.7	45.9	12.4
2002	10.4	47.5	11.8
2003	10.1	51.3	11.5
2004	10.9	49.6	10.9
2005	10.4	50.8	10.9
2006	10.4	50.6	10.7
2007	9.8	51.2	10.5
2008	10.2	50.5	10.7
2009	10.4	52.6	10.7
2010	10.5	52.5	10.4
2011	9.9	51.4	10.9
2012	9.9	51.8	10.6
2013	9.9	53.1	10.6
2014	10.1	52.4	10.7
2015	10.2	54.0	10.4
2016	9.7	55.8	10.3

¹ Rate of very low birthweight (less than 1,500 grams) per 1,000 live births.

² Rate of low birthweight (1,500 - 2,499 grams) per 1,000 live births.

³ Percent of macrosomia (birthweight greater than 4,000 grams) per 100 live births.

Rates and percentages are calculated excluding missing and unknown values.