

Induced abortions by length of gestation, procedure type, complications, and age of patient, Oregon residents, 2011-2013

State Total

Medical and health characteristics	Total	Patient's age				
		10-14	15-17	18-19	20+	N.S.
Total	25,056	70	1,029	2,132	21,640	185
Gestation (weeks)						
< 9	16,155	29	593	1,294	14,120	119
9-12	5,353	20	265	497	4,550	21
13-16	1,714	7	96	168	1,430	13
17-20	791	4	36	86	660	5
21-22	269	4	14	19	231	1
23+	147	2	5	15	125	—
Unknown	627	4	20	53	524	26
Contraceptive use						
None	17,499	51	766	1,573	14,990	119
Birth control pills	2,160	—	60	203	1,876	21
Condoms	3,390	16	135	240	2,974	25
Other contraceptive	1,590	—	30	72	1,468	20
Unknown	718	3	46	70	593	6
Procedure type						
Suction curette	12,165	36	545	1,050	10,458	76
Medical (non-surgical)	6,855	13	251	598	5,920	73
Dilation & evacuation	5,837	21	228	470	5,083	35
Intra-uterine instillation	22	—	1	4	17	—
Vaginal prostaglandin	43	—	—	1	41	1
Sharp curettage	4	—	—	—	4	—
Other	99	—	2	8	89	—
Unknown	31	—	2	1	28	—
Complications						
None	24,386	69	1,008	2,081	21,058	170
Hemorrhage	26	1	1	1	21	2
Infection	49	—	3	4	40	2
Uterine perforation	2	—	—	—	2	—
Cervical laceration	3	—	—	1	2	—
Retained products	271	—	4	20	244	3
Failure of first method	81	—	1	8	69	3
Other	167	—	12	16	136	3
Multiple complications	67	—	—	1	64	2
Unknown	4	—	—	—	4	—

— Quantity is zero.

NOTE: Includes reported abortions to Oregon residents including those obtained out-of-state. Because some states (e.g., California) do not record data on residence for abortion patients, not all out-of-state abortions are included.