

Induced abortions by length of gestation, procedure type, complications, and age of patient, Oregon residents, 2012-2014

State total

Medical and health characteristics	Total	Patient's age				
		10-14	15-17	18-19	20+	N.S.
Total	23,825	68	916	1,850	20,664	327
Gestation (weeks)						
< 9	15,269	31	505	1,109	13,428	196
9-12	5,122	20	250	465	4,320	67
13-16	1,631	5	80	147	1,377	22
17-20	775	4	36	67	660	8
21-22	279	3	13	15	245	3
23+	136	1	9	12	113	1
Unknown	613	4	23	35	521	30
Contraceptive use						
None	16,635	51	706	1,364	14,282	232
Birth control pills	2,000	2	50	179	1,741	28
Condoms	3,127	13	112	191	2,771	40
Other contraceptive	1,636	—	23	72	1,513	28
Unknown	728	2	33	67	612	14
Procedure type						
Suction curette	10,932	38	446	884	9,424	140
Medical (non-surgical)	6,964	13	239	550	6,053	109
Dilation & evacuation	5,761	17	227	405	5,038	74
Intra-uterine instillation	27	—	2	4	20	1
Vaginal prostaglandin	32	—	—	2	29	1
Sharp curettage	6	—	1	—	5	—
Other	74	—	1	5	68	—
Unknown	29	—	—	—	27	2
Complications						
None	23,177	67	896	1,805	20,099	310
Hemorrhage	17	1	—	—	15	1
Infection	36	—	2	4	28	2
Uterine perforation	3	—	—	—	3	—
Cervical laceration	2	—	—	1	1	—
Retained products	206	—	4	12	186	4
Failure of first method	59	—	1	4	52	2
Other	238	—	12	22	198	6
Multiple complications	81	—	1	2	76	2
Unknown	6	—	—	—	6	—

— Quantity is zero.

NOTE: Includes reported abortions to Oregon residents including those obtained out-of-state. Because some states (e.g., California) do not record data on residence for abortion patients, not all out-of-state abortions are included.