

Antibiotics are drugs like penicillin that are used to fight or prevent bacterial infection. In the past 4 weeks, have you taken any antibiotics?

(Weighted Column Percents)

	Total	Age Groups					
		18-24	25-34	35-44	45-54	55-64	65+
Yes	9.3	6.3	10.2	9.1	8.4	11.1	10.2
No	90.7	93.7	89.8	90.9	91.6	88.9	89.8
Male							
Yes	7.2	5.7	10.1	7.1	4.8	8.2	7.5
No	92.8	94.3	89.9	92.9	95.2	91.8	92.5
Female							
Yes	11.2	6.9	10.3	11.2	12.0	13.8	12.2
No	88.8	93.1	89.7	88.8	88.0	86.2	87.8

Denominator is:
All survey respondents.

Antibiotics are drugs like penicillin that are used to fight or prevent bacterial infection. In the past 4 weeks, have you taken any antibiotics?

(Unweighted Frequencies)

	Total	Age Groups					
		18-24	25-34	35-44	45-54	55-64	65+
Total	3,096	233	483	549	666	538	627
Yes	312	21	48	51	64	61	67
No	2,769	212	433	498	601	474	551
Don't know/Not sure	14		2		1	3	8
Refused	1						1
Male							
Total	1,237	96	206	206	277	220	232
Yes	99	8	20	15	16	20	20
No	1,133	88	186	191	261	199	208
Don't know/Not sure	4					1	3
Refused	1						1
Female							
Total	1,859	137	277	343	389	318	395
Yes	213	13	28	36	48	41	47
No	1,636	124	247	307	340	275	343
Don't know/Not sure	10		2		1	2	5

Denominator is:
All survey respondents.

How much do you agree/disagree with the following statement?: I will be sick for a longer time if I don't receive an antibiotic for cough, cold, or flu symptoms.

(Weighted Column Percents)

		Total	Age Groups					
			18-24	25-34	35-44	45-54	55-64	65+
Strongly agree		18.7	21.5	18.5	15.2	16.3	22.4	20.6
Somewhat agree		15.5	26.7	14.2	12.3	12.1	13.9	17.1
Neither agree or disagree		1.4		1.7	1.6	1.0	1.2	2.7
Somewhat disagree		25.5	29.5	27.4	25.6	24.4	19.9	26.0
Strongly disagree		39.0	22.4	38.3	45.3	46.3	42.6	33.6
Male	Strongly agree	19.3	23.2	21.9	13.9	18.3	18.6	22.0
	Somewhat agree	17.0	28.8	15.7	13.4	14.0	17.0	16.4
	Neither agree or disagree	1.3		1.3	2.1	.8	1.2	2.5
	Somewhat disagree	26.5	28.1	25.5	30.9	26.3	21.4	25.2
	Strongly disagree	35.8	19.8	35.6	39.8	40.5	41.8	33.8
Female	Strongly agree	18.1	19.6	15.0	16.7	14.4	26.0	19.6
	Somewhat agree	14.0	24.4	12.7	11.2	10.1	11.0	17.6
	Neither agree or disagree	1.4		2.0	1.0	1.1	1.2	2.9
	Somewhat disagree	24.4	30.9	29.3	20.1	22.4	18.6	26.5
	Strongly disagree	42.0	25.1	41.0	51.0	52.0	43.2	33.5

Denominator is:
All survey respondents.

How much do you agree/disagree with the following statement?: I will be sick for a longer time if I don't receive an antibiotic for cough, cold, or flu symptoms.

(Unweighted Frequencies)

		Total	Age Groups					
			18-24	25-34	35-44	45-54	55-64	65+
Total		3,096	233	483	549	666	538	627
Strongly agree		530	45	80	85	98	115	107
Somewhat agree		421	57	70	61	73	67	93
Neither agree or disagree		39		6	8	7	6	12
Somewhat disagree		738	75	132	133	144	108	146
Strongly disagree		1,213	54	180	248	322	219	190
Don't know/Not sure		140	2	14	13	17	22	72
Refused		15		1	1	5	1	7
Male	Total	1,237	96	206	206	277	220	232
	Strongly agree	218	20	39	31	46	38	44
	Somewhat agree	190	24	32	24	39	36	35
	Neither agree or disagree	16		2	4	2	3	5
	Somewhat disagree	301	29	53	62	63	45	49
	Strongly disagree	448	21	70	80	119	87	71
	Don't know/Not sure	57	2	9	5	5	10	26
	Refused	7		1		3	1	2
Female	Total	1,859	137	277	343	389	318	395
	Strongly agree	312	25	41	54	52	77	63
	Somewhat agree	231	33	38	37	34	31	58
	Neither agree or disagree	23		4	4	5	3	7
	Somewhat disagree	437	46	79	71	81	63	97
	Strongly disagree	765	33	110	168	203	132	119
	Don't know/Not sure	83		5	8	12	12	46
	Refused	8			1	2		5

Denominator is:
All survey respondents.

How much do you agree/disagree with the following statement?: By the time I am sick enough to talk to or visit a doctor because of a cold, I usually expect a prescription for antibiotics.

(Weighted Column Percents)

	Total	Age Groups					
		18-24	25-34	35-44	45-54	55-64	65+
Strongly agree	25.2	27.7	26.4	28.4	24.2	25.6	19.1
Somewhat agree	16.4	27.7	16.1	15.2	12.0	13.9	16.8
Neither agree or disagree	1.6	1.2	2.0	.8	1.2	.2	4.2
Somewhat disagree	20.4	25.3	24.1	18.7	17.9	17.3	19.9
Strongly disagree	36.3	18.1	31.4	36.9	44.7	42.9	40.1
Male							
Strongly agree	24.4	24.1	27.7	28.2	22.8	24.3	16.9
Somewhat agree	18.5	32.6	21.8	17.6	10.0	14.2	17.7
Neither agree or disagree	1.8	2.2	1.9	.3	1.1	.5	5.4
Somewhat disagree	20.8	20.5	22.8	19.7	23.4	17.9	19.2
Strongly disagree	34.6	20.5	25.8	34.1	42.7	43.1	40.7
Female							
Strongly agree	26.1	31.6	25.1	28.5	25.6	26.9	20.6
Somewhat agree	14.5	22.4	10.2	12.6	13.9	13.6	16.1
Neither agree or disagree	1.5		2.1	1.4	1.3		3.2
Somewhat disagree	20.0	30.6	25.5	17.7	12.5	16.8	20.5
Strongly disagree	38.0	15.4	37.0	39.8	46.6	42.8	39.6

Denominator is:
All survey respondents.

How much do you agree/disagree with the following statement?: By the time I am sick enough to talk to or visit a doctor because of a cold, I usually expect a prescription for antibiotics.

(Unweighted Frequencies)

		Total	Age Groups					
			18-24	25-34	35-44	45-54	55-64	65+
Total		3,096	233	483	549	666	538	627
Strongly agree		717	67	122	143	147	132	106
Somewhat agree		444	61	71	77	75	68	92
Neither agree or disagree		44	2	8	6	7	2	19
Somewhat disagree		592	60	111	105	116	85	115
Strongly disagree		1,162	41	154	207	302	230	228
Don't know/Not sure		120	2	16	10	14	21	57
Refused		17		1	1	5		10
Male	Total	1,237	96	206	206	277	220	232
	Strongly agree	268	21	51	56	54	50	36
	Somewhat agree	201	32	42	37	26	28	36
	Neither agree or disagree	20	2	3	1	3	2	9
	Somewhat disagree	244	20	43	39	66	37	39
	Strongly disagree	444	20	55	70	122	93	84
	Don't know/Not sure	55	1	11	3	6	10	24
	Refused	5		1				4
Female	Total	1,859	137	277	343	389	318	395
	Strongly agree	449	46	71	87	93	82	70
	Somewhat agree	243	29	29	40	49	40	56
	Neither agree or disagree	24		5	5	4		10
	Somewhat disagree	348	40	68	66	50	48	76
	Strongly disagree	718	21	99	137	180	137	144
	Don't know/Not sure	65	1	5	7	8	11	33
	Refused	12			1	5		6

Denominator is:
All survey respondents.

How often are antibiotics needed for a sore throat?

(Weighted Column Percents)

		Total	Age Groups					
			18-24	25-34	35-44	45-54	55-64	65+
Always		2.6	3.8	3.0	2.0	1.7	2.6	2.9
Almost always		4.7	5.7	4.2	4.9	4.4	4.5	4.8
Sometimes		44.8	52.4	43.7	40.6	47.8	45.1	41.2
Almost never		31.3	25.9	33.3	35.7	30.9	31.9	28.6
Never		16.5	12.3	15.7	16.8	15.2	15.9	22.5
Male	Always	3.2	5.5	4.3	2.1	1.4	2.9	4.1
	Almost always	4.1	1.7	4.4	6.1	4.5	3.6	3.4
	Sometimes	41.8	52.8	39.1	37.7	45.2	41.0	36.1
	Almost never	32.4	27.7	36.2	35.1	30.9	30.9	31.8
	Never	18.5	12.4	16.0	19.0	18.0	21.7	24.6
Female	Always	2.0	1.9	1.7	1.9	1.9	2.4	2.2
	Almost always	5.2	10.1	3.9	3.7	4.2	5.3	5.8
	Sometimes	47.8	51.9	48.6	43.6	50.5	49.0	44.8
	Almost never	30.4	24.0	30.4	36.2	31.0	33.0	26.3
	Never	14.6	12.1	15.4	14.6	12.5	10.3	20.9

Denominator is:
All survey respondents.

How often are antibiotics needed for a sore throat?

(Unweighted Frequencies)

		Total	Age Groups					65+
			18-24	25-34	35-44	45-54	55-64	
Total		3,096	233	483	549	666	538	627
Always		62	6	11	8	8	13	16
Almost always		133	13	19	25	24	21	31
Sometimes		1,347	119	218	222	308	238	242
Almost never		995	66	164	196	221	176	172
Never		488	26	66	88	95	79	134
Don't know/Not sure		62	3	4	9	9	9	28
Refused		9		1	1	1	2	4
Male	Total	1,237	96	206	206	277	220	232
	Always	30	3	7	3	3	6	8
	Almost always	49	2	9	13	11	7	7
	Sometimes	489	50	80	76	118	89	76
	Almost never	409	26	77	73	92	70	71
	Never	227	13	29	37	49	44	55
	Don't know/Not sure	30	2	4	4	3	4	13
	Refused	3				1		2
Female	Total	1,859	137	277	343	389	318	395
	Always	32	3	4	5	5	7	8
	Almost always	84	11	10	12	13	14	24
	Sometimes	858	69	138	146	190	149	166
	Almost never	586	40	87	123	129	106	101
	Never	261	13	37	51	46	35	79
	Don't know/Not sure	32	1		5	6	5	15
	Refused	6		1	1		2	2

Denominator is:
All survey respondents.

How often are antibiotics needed for a cold?

(Weighted Column Percents)

		Total	Age Groups					
			18-24	25-34	35-44	45-54	55-64	65+
Always		2.3	4.3	3.4	2.2	1.2	1.5	1.4
Almost always		2.4	4.7	1.8	2.0	2.3	1.4	2.8
Sometimes		21.9	34.8	23.4	21.2	15.7	20.5	19.7
Almost never		26.0	29.4	25.5	23.6	28.0	24.1	25.5
Never		47.4	26.9	45.9	51.0	52.9	52.4	50.6
Male	Always	2.7	5.1	4.6	2.6	2.0	1.0	.6
	Almost always	2.7	7.0	.4	2.4	1.9	2.1	3.4
	Sometimes	25.1	35.2	29.7	25.2	18.4	22.6	20.8
	Almost never	27.6	32.2	26.7	24.4	32.1	22.1	27.3
	Never	42.0	20.4	38.5	45.4	45.7	52.2	47.9
Female	Always	1.9	3.3	2.2	1.9	.5	2.1	2.0
	Almost always	2.1	2.1	3.2	1.5	2.7	.7	2.4
	Sometimes	18.9	34.2	16.8	17.1	13.0	18.4	18.9
	Almost never	24.4	26.4	24.1	22.9	23.9	26.1	24.2
	Never	52.7	34.0	53.7	56.6	60.0	52.7	52.6

Denominator is:
All survey respondents.

How often are antibiotics needed for a cold?

(Unweighted Frequencies)

		Total	Age Groups					65+
			18-24	25-34	35-44	45-54	55-64	
Total		3,096	233	483	549	666	538	627
Always		50	6	14	7	7	9	7
Almost always		61	10	7	10	10	7	17
Sometimes		587	81	103	95	97	98	113
Almost never		805	73	124	137	177	138	156
Never		1,540	62	231	294	367	276	310
Don't know/Not sure		47	1	3	5	7	9	22
Refused		6		1	1	1	1	2
Male	Total	1,237	96	206	206	277	220	232
	Always	24	4	8	4	5	2	1
	Almost always	27	6	1	5	4	4	7
	Sometimes	280	36	58	46	48	47	45
	Almost never	340	31	57	53	86	51	62
	Never	539	19	79	94	130	112	105
	Don't know/Not sure	24		3	3	3	4	11
	Refused	3			1	1		1
Female	Total	1,859	137	277	343	389	318	395
	Always	26	2	6	3	2	7	6
	Almost always	34	4	6	5	6	3	10
	Sometimes	307	45	45	49	49	51	68
	Almost never	465	42	67	84	91	87	94
	Never	1,001	43	152	200	237	164	205
	Don't know/Not sure	23	1		2	4	5	11
	Refused	3		1			1	1

Denominator is:
All survey respondents.

How often are antibiotics needed for a cough without fever?

(Weighted Column Percents)

		Total	Age Groups					65+
			18-24	25-34	35-44	45-54	55-64	
Always		2.2	2.8	3.0	3.1	.9	1.9	1.7
Almost always		2.3	6.0	1.9	1.7	.9	3.0	1.6
Sometimes		19.0	29.0	22.3	16.7	16.8	14.9	16.3
Almost never		27.3	30.8	31.1	25.9	28.8	25.2	22.4
Never		46.1	30.3	40.0	50.0	50.1	52.5	50.1
Don't know/Not sure		2.9	1.1	1.5	2.4	2.1	2.5	7.6
Refused		.2		.1	.2	.3		.3
Male	Always	2.2	3.5	2.9	3.6	1.6	.9	.5
	Almost always	3.1	10.5	2.3	1.5	.9	3.3	2.4
	Sometimes	18.5	28.7	22.9	15.7	18.1	13.9	12.1
	Almost never	27.4	29.6	30.6	26.3	30.3	23.1	22.7
	Never	44.9	26.0	38.7	49.2	46.8	54.6	52.6
	Don't know/Not sure	3.6	1.7	2.6	3.3	1.7	4.1	9.3
	Refused	.3			.5	.6		.4
Female	Always	2.2	2.0	3.1	2.6	.2	2.8	2.5
	Almost always	1.5	1.1	1.4	1.8	1.0	2.7	1.0
	Sometimes	19.4	29.3	21.7	17.8	15.5	15.9	19.3
	Almost never	27.3	32.0	31.6	25.6	27.4	27.3	22.2
	Never	47.3	35.0	41.5	50.8	53.4	50.5	48.3
	Don't know/Not sure	2.3	.5	.3	1.5	2.5	.9	6.4
	Refused	.1		.3				.3

Denominator is:
All survey respondents.

How often are antibiotics needed for a cough without fever?

(Unweighted Frequencies)

		Total	Age Groups					65+
			18-24	25-34	35-44	45-54	55-64	
Total		3,096	233	483	549	666	538	627
Always		48	5	11	9	4	9	10
Almost always		53	10	8	9	3	13	10
Sometimes		559	66	104	93	110	86	100
Almost never		865	78	158	148	200	146	135
Never		1,474	72	196	276	332	271	327
Don't know/Not sure		92	2	5	13	16	13	43
Refused		5		1	1	1		2
Male	Total	1,237	96	206	206	277	220	232
	Always	18	3	4	5	3	2	1
	Almost always	30	8	5	4	2	6	5
	Sometimes	210	27	44	32	47	32	28
	Almost never	348	30	68	56	89	56	49
	Never	582	27	81	101	130	115	128
	Don't know/Not sure	46	1	4	7	5	9	20
	Refused	3			1	1		1
Female	Total	1,859	137	277	343	389	318	395
	Always	30	2	7	4	1	7	9
	Almost always	23	2	3	5	1	7	5
	Sometimes	349	39	60	61	63	54	72
	Almost never	517	48	90	92	111	90	86
	Never	892	45	115	175	202	156	199
	Don't know/Not sure	46	1	1	6	11	4	23
	Refused	2		1				1

Denominator is:
All survey respondents.

How often are antibiotics needed for a runny nose with yellow or green mucous?

(Weighted Column Percents)

		Total	Age Groups					
			18-24	25-34	35-44	45-54	55-64	65+
Always		9.9	9.0	13.0	11.7	11.0	7.3	6.2
Almost always		16.1	24.7	14.9	15.9	14.7	17.1	11.6
Sometimes		39.7	37.1	39.3	44.9	41.7	40.6	32.9
Almost never		16.9	16.5	17.9	13.4	20.4	16.5	16.1
Never		17.4	12.7	14.9	14.0	12.1	18.6	33.3
Male	Always	8.1	4.9	11.9	10.9	10.1	3.6	4.1
	Almost always	14.0	24.2	13.4	13.9	10.6	14.2	9.6
	Sometimes	36.3	38.1	35.7	38.3	40.3	35.0	27.6
	Almost never	18.6	15.0	19.8	17.1	21.4	20.7	16.9
	Never	22.9	17.8	19.3	19.8	17.6	26.4	41.8
Female	Always	11.6	13.5	14.2	12.6	12.0	10.6	7.6
	Almost always	18.0	25.3	16.5	17.9	18.6	19.7	13.0
	Sometimes	42.9	36.0	43.0	51.6	43.2	45.7	36.5
	Almost never	15.2	18.1	15.9	9.7	19.5	12.6	15.5
	Never	12.2	7.2	10.4	8.2	6.8	11.4	27.4

Denominator is:
All survey respondents.

How often are antibiotics needed for a runny nose with yellow or green mucous?

(Unweighted Frequencies)

		Total	Age Groups					65+
			18-24	25-34	35-44	45-54	55-64	
Total		3,096	233	483	549	666	538	627
Always		283	20	56	62	70	38	37
Almost always		462	54	77	86	94	87	64
Sometimes		1,218	90	190	248	278	211	201
Almost never		483	37	84	70	124	80	88
Never		511	27	63	69	77	90	185
Don't know/Not sure		133	4	12	13	23	32	49
Refused		6	1	1	1			3
Male	Total	1,237	96	206	206	277	220	232
	Always	88	4	22	20	25	8	9
	Almost always	149	17	29	27	30	28	18
	Sometimes	423	43	71	75	105	71	58
	Almost never	227	14	40	38	56	42	37
	Never	276	16	36	38	49	52	85
	Don't know/Not sure	71	1	8	7	12	19	24
	Refused	3	1		1			1
Female	Total	1,859	137	277	343	389	318	395
	Always	195	16	34	42	45	30	28
	Almost always	313	37	48	59	64	59	46
	Sometimes	795	47	119	173	173	140	143
	Almost never	256	23	44	32	68	38	51
	Never	235	11	27	31	28	38	100
	Don't know/Not sure	62	3	4	6	11	13	25
	Refused	3		1				2

Denominator is:
All survey respondents.

Are you aware of any dangers to yourself or other people associated with taking antibiotics?

(Weighted Column Percents)

		Total	Age Groups					65+
			18-24	25-34	35-44	45-54	55-64	
Yes		44.9	34.2	46.2	50.0	56.8	47.9	29.8
No		53.5	63.9	52.7	48.5	42.4	51.7	66.0
Don't know/Not sure		1.5	1.9	.8	1.2	.8	.4	4.2
Refused		.1		.3	.2			.1
Male	Yes	40.4	28.9	41.8	44.6	52.8	37.9	29.2
	No	57.8	68.7	56.7	54.3	46.5	61.9	65.3
	Don't know/Not sure	1.7	2.4	1.0	.7	.7	.2	5.5
	Refused	.2		.4	.5			
Female	Yes	49.2	40.1	50.8	55.6	60.7	57.6	30.2
	No	49.3	58.6	48.4	42.6	38.4	41.9	66.5
	Don't know/Not sure	1.4	1.4	.5	1.8	.9	.5	3.2
	Refused	.1		.3				.1

Denominator is:
All survey respondents.

Are you aware of any dangers to yourself or other people associated with taking antibiotics?

(Unweighted Frequencies)

		Total	Age Groups					65+
			18-24	25-34	35-44	45-54	55-64	
Total		3,096	233	483	549	666	538	627
Yes		1,447	85	233	289	384	263	193
No		1,606	144	246	255	277	273	411
Don't know/Not sure		39	4	2	4	5	2	22
Refused		4		2	1			1
Male	Total	1,237	96	206	206	277	220	232
	Yes	510	26	89	97	146	82	70
	No	708	68	115	107	129	137	152
	Don't know/Not sure	17	2	1	1	2	1	10
	Refused	2		1	1			
Female	Total	1,859	137	277	343	389	318	395
	Yes	937	59	144	192	238	181	123
	No	898	76	131	148	148	136	259
	Don't know/Not sure	22	2	1	3	3	1	12
	Refused	2		1				1

Denominator is:
All survey respondents.

What kind of health dangers do you see to yourself and other people associated with taking antibiotics?

(Weighted Column Percents)

		Total	Age Groups					
			18-24	25-34	35-44	45-54	55-64	65+
Male	Not aware of any dangers	55.1	65.8	53.7	49.9	43.2	52.1	70.2
	Emerging drug resistance	25.1	12.6	23.7	30.9	35.6	29.6	14.2
	Allergies or reactions to antibiotics	17.5	12.3	18.6	18.1	23.9	17.3	12.3
	Antibiotics likely to kill friendly bacteria	5.7	4.2	5.9	5.8	7.9	8.1	2.3
	Unhealthy practice to take drugs or chemicals in general	1.5	.8	1.3	1.6	1.2	2.0	2.2
	Other reason	8.5	7.5	9.3	9.9	10.7	8.2	4.8
	Aware of danger, but gave no reason(s)	3.0	5.1	3.0	3.4	2.6	2.0	2.2
	Not aware of any dangers	59.6	71.1	58.0	55.2	47.2	62.1	70.8
	Emerging drug resistance	22.0	8.6	18.9	28.5	30.7	24.7	15.6
	Allergies or reactions to antibiotics	14.4	11.6	18.1	13.8	20.7	7.8	10.5
Female	Antibiotics likely to kill friendly bacteria	5.2	5.2	5.5	3.5	8.1	7.0	1.9
	Unhealthy practice to take drugs or chemicals in general	1.1	.7	.3	.9	1.8	1.1	1.7
	Other reason	7.7	4.2	8.5	9.7	10.2	7.5	4.3
	Aware of danger, but gave no reason(s)	2.8	3.3	2.5	2.2	3.4	3.2	2.5
	Not aware of any dangers	50.8	59.9	49.0	44.4	39.3	42.4	69.8
	Emerging drug resistance	28.2	16.9	28.8	33.3	40.4	34.3	13.3
	Allergies or reactions to antibiotics	20.5	13.1	19.2	22.4	27.0	26.6	13.6
	Antibiotics likely to kill friendly bacteria	6.2	3.1	6.3	8.3	7.7	9.1	2.7
	Unhealthy practice to take drugs or chemicals in general	2.0	.8	2.4	2.2	.7	2.9	2.5
	Other reason	9.3	11.2	10.2	10.2	11.1	8.8	5.1
Aware of danger, but gave no reason(s)	3.1	7.0	3.5	4.6	1.7	.9	1.9	

Denominator is:

All survey respondents who said they were aware of any dangers associated with taking antibiotics.

What kind of health dangers do you see to yourself and other people associated with taking antibiotics?

		Total	Age Groups					
			18-24	25-34	35-44	45-54	55-64	65+
Total		3,096	233	483	549	666	538	627
	Not aware of any dangers	1,645	148	248	259	282	275	433
	Emerging drug resistance	853	36	125	182	253	164	93
	Allergies or reactions to antibiotics	567	34	94	103	157	101	78
	Antibiotics likely to kill friendly bacteria	193	11	31	37	56	43	15
	Unhealthy practice to take drugs or chemicals in general	53	2	7	10	9	12	13
	Other reason	271	20	48	56	71	43	33
	Aware of danger, but gave no reason(s)	81	10	14	18	14	10	15
	Refused	12	1	3	4	1	1	2
Male	Not aware of any dangers	725	70	116	108	131	138	162
	Emerging drug resistance	301	8	44	64	93	55	37
	Allergies or reactions to antibiotics	173	11	37	27	52	20	26
	Antibiotics likely to kill friendly bacteria	69	6	11	7	25	15	5
	Unhealthy practice to take drugs or chemicals in general	16	1	1	2	5	3	4
	Other reason	93	4	17	19	29	14	10
	Aware of danger, but gave no reason(s)	31	2	6	6	6	6	5
	Refused	8		2	3	1	1	1
Female	Not aware of any dangers	920	78	132	151	151	137	271
	Emerging drug resistance	552	28	81	118	160	109	56
	Allergies or reactions to antibiotics	394	23	57	76	105	81	52
	Antibiotics likely to kill friendly bacteria	124	5	20	30	31	28	10
	Unhealthy practice to take drugs or chemicals in general	37	1	6	8	4	9	9
	Other reason	178	16	31	37	42	29	23
	Aware of danger, but gave no reason(s)	50	8	8	12	8	4	10
	Refused	4	1	1	1			1

Denominator is:

All survey respondents who said they were aware of any dangers associated with taking antibiotics.