

## Tobacco use and related topics among adults by race and ethnicity, Oregon 2010-2011



Use this table to estimate the overall burden of disease within each race and ethnicity

	White, NL		African American, NL		Asian or Pacific Islander, NL		American Indian or Alaska Native, NL		Latino	
	Unadjusted (%)	Number of adults	Unadjusted (%)	Number of adults	Unadjusted (%)	Number of adults	Unadjusted (%)	Number of adults	Unadjusted (%)	Number of adults
<b>Cigarettes</b>										
Current smoker (female)	18.5	236,500	24.7	6,300	11.9 ^	8,500	33.8	6,700	15.7	20,400
Current smoker (male)	21.8	263,700	41.4	12,600	18.0 ^	10,700	35.8	6,800	30.0	44,200
Current smoker (total)	20.1	500,000	33.9	19,000	14.7	19,300	34.7	13,500	23.2	64,300
Smokes some days	5.1	126,900	9.7	5,400	6.6 ^	8,600	9.6	3,700	8.3	23,000
Smokes every day	15.0	373,200	24.2	13,500	8.1 ^	10,600	25.1	9,800	14.9	41,300
Started smoking regularly before age 18	53.4	1,328,500	33.2	18,600	39.4	51,600	59.4	23,100	60.7	168,100
<b>Quitting among current cigarette smokers</b>										
Preparing to quit	20.9	104,500	42.2	8,000	--	--	34.5	4,700	30.9 ^	19,900
Attempted to quit during previous year	51.9	259,500	68.6	13,000	63.9	12,300	47.9	6,500	65.7	42,200
Has heard about the Oregon Tobacco Quit Line	44.2	1,099,600	34.8	19,500	32.6	42,700	48.1	18,700	34.0	94,200
<b>Smokeless tobacco</b>										
Smokeless tobacco use (male)	8.5	211,500	--	--	0.9 ^	1,200	11.8	4,600	4.8 ^	13,300

NL = non-Latino

^ This number may be statistically unreliable and should be interpreted with caution.

-- This number is suppressed because it is statistically unreliable.

## Tobacco use and related topics among adults by race and ethnicity, Oregon 2010-2011 (continued from page 1)

Use this table to estimate the overall burden of disease within each race and ethnicity

	White, NL		African American, NL		Asian or Pacific Islander, NL		American Indian or Alaska Native, NL		Latino	
	Unadjusted (%)	Number of adults	Unadjusted (%)	Number of adults	Unadjusted (%)	Number of adults	Unadjusted (%)	Number of adults	Unadjusted (%)	Number of adults
<b>Secondhand smoke</b>										
Smoking is never allowed in the home	91.0	2,263,900	79.5	44,500	92.7	121,400	85.6	33,300	91.6	253,700
Smoking inside the home in last 30 days	6.5	161,700	14.1	7,900	2.5 ^	3,300	13.2	5,100	4.7 ^	13,000
Exposed to secondhand smoke indoors	20.8	517,500	27.6	15,400	17.1	22,400	39.8	15,500	17.7	49,000
Smoking is never allowed in the car	79.0	1,965,300	63.5	35,500	91.1	119,300	68.9	26,800	89.3	247,400
<b>Tobacco health services</b>										
HCP asked about tobacco use (everyone)	64.7	1,609,600	74.0	41,400	61.2	80,200	79.9	31,100	65.9	182,500
HCP advised to quit tobacco use (current smokers)	57.3	286,500	61.2	11,600	63.0	12,200	47.1	6,400	49.8	32,000
HCP offered assistance to quit tobacco use (current smokers)	39.2	196,000	48.3	9,200	41.4 ^	8,000	39.0	5,300	13.8 ^	8,900

NL = non-Latino; HCP = Health care provider

^ This number may be statistically unreliable and should be interpreted with caution.

## Tobacco use and related topics among adults by race and ethnicity, Oregon 2010-2011

Use this table to compare burden of disease among race and ethnicity groups

	Age-adjusted (%)				
	White, NL	African American, NL	Asian or Pacific Islander, NL	American Indian or Alaska Native, NL	Latino
<b>Cigarettes</b>					
Current smoker (female)	20.2	24.3	11.3 ^	33.9	14.5
Current smoker (male)	22.8	40.7	18.0 ^	36.9	26.5
Current smoker (total)	21.4	33.3	14.1	35.3	20.8
Smokes some days	5.5	9.4	6.0 ^	9.7	7.7
Smokes every day	16.0	23.9	8.2 ^	25.6	13.1
Started smoking regularly before age 18	55.3	34.2	41.9	58.4	59.9
<b>Quitting among current cigarette smokers</b>					
Preparing to quit	20.6	41.3	--	34.6	30.7 ^
Attempted to quit during previous year	51.2	66.0	54.9	48.7	54.6
Has heard about the Oregon Tobacco Quit Line	44.4	36.3	33.7	49.0	33.4
<b>Smokeless tobacco</b>					
Smokeless tobacco use (male)	9.3	--	0.9 ^	11.7	3.8 ^

NL = non-Latino

^ This number may be statistically unreliable and should be interpreted with caution.

-- This number is suppressed because it is statistically unreliable.

## Tobacco use and related topics among adults by race and ethnicity, Oregon 2010-2011 (continued from page 3)

Use this table to compare burden of disease among race and ethnicity groups

	Age-adjusted (%)				
	White, NL	African American, NL	Asian or Pacific Islander, NL	American Indian or Alaska Native, NL	Latino
<b>Secondhand smoke</b>					
Smoking is never allowed in the home	91.1	80.0	92.5	84.5	89.0
Smoking inside the home in last 30 days	6.7	14.0	2.6 ^	14.1	4.6 ^
One or more hours per week in same room with smoking	22.1	27.3	15.9	40.1	18.7
Smoking is never allowed in the car	77.9	65.4	90.6	68.8	90.5
<b>Tobacco health services</b>					
HCP asked about tobacco use (everyone)	65.8	74.3	60.5	81.0	64.0
HCP advised to quit tobacco use (current smokers)	58.2	65.5	68.7	49.1	45.2
HCP offered assistance to quit tobacco use (current smokers)	40.3	51.1	53.3 ^	40.1	29.0 ^

NL = non-Latino; HCP = Health care provider

^ This number may be statistically unreliable and should be interpreted with caution.

Source: Oregon Behavioral Risk Factors Surveillance System Race Oversample Dataset 2010-2011; age-adjusted to the 2000 standard population.

Note: For an explanation of using unadjusted versus age-adjusted estimates, please see <http://public.health.oregon.gov/DiseasesConditions/ChronicDisease/DataReports/Pages/TechnicalNotes.aspx>.

Suggested citation: Oregon Health Authority, Public Health Division, Health Promotion and Chronic Disease Prevention section. Tobacco use and related topics among adults by race and ethnicity, Oregon 2010-2011. <http://public.health.oregon.gov/DiseasesConditions/ChronicDisease/DataReports/Pages/AdultData.aspx>.

Created August 11, 2014. Accessed [DOWNLOAD DATE].