



## HEALTH HARMS FROM SMOKING AND OTHER TOBACCO USE

---

Tobacco use kills more than 400,000 people each year in the United States, or more than the total number killed by AIDS, alcohol, motor vehicles, homicide, illegal drugs, and suicide combined.<sup>1</sup> Among current smokers 57 percent of all male deaths and nearly 50 percent of all deaths in women are attributed to smoking.<sup>2</sup> Even if the number of smoking related deaths were cut in half, smoking would still kill more people than all of these other causes.

In 1964, the Surgeon General first documented the harmful effects of smoking in *Smoking and Health: Report of the Advisory Committee of the Surgeon General of the Public Health Service*, which summarized the state of the science knowledge regarding tobacco use at that time.<sup>3</sup> Research conducted since then has firmly established that smoking and other forms of tobacco consumption cause an enormous amount of health problems and related death and suffering. Today, smoking is the leading preventable cause of death in the United States. Despite the numerous reports of the Surgeon General and the National Institute for Health on the risks of smoking, 46 million Americans still smoke,<sup>4</sup> and approximately half of all continuing smokers will die prematurely as a result of their habit.

Although most of the research on health harms focuses on cigarette smoking, recent studies have shown that other tobacco use, such as cigar and pipe smoking, also produce similar health risks to users.<sup>5</sup>

**Smoking-Caused Cancer.** Smoking is responsible for 87 percent of lung cancer deaths (90 percent in men, 80 percent in women).<sup>6</sup> Over 125,000 men and women die of smoking caused lung cancer each year.<sup>7</sup> Compared to nonsmokers, men who smoke are about 23 times more likely to develop lung cancer and women who smoke are about 13 times more likely.<sup>8</sup> Beyond just lung cancer, thirty percent of all cancer deaths are due to smoking.<sup>9</sup> Smoking is a known cause of cancer of the lung, larynx, oral cavity, bladder, pancreas, uterus, cervix, kidney, stomach and esophagus.<sup>10</sup>

**Smoking-Caused Respiratory Diseases.** Twenty-three percent of smoking-attributable deaths, or more than 100,000 smoking deaths per year, involve respiratory diseases.<sup>11</sup> Smoking is a known cause of most cases of chronic obstructive pulmonary disease (COPD) which includes emphysema and chronic bronchitis.<sup>12</sup> Smoking is accountable for more than 90 percent of all COPD deaths.<sup>13</sup> Male and female smokers increase their risk of death from bronchitis and emphysema by 10 times.<sup>14</sup>

**Smoking-Caused Heart Disease and Heart Attacks.** Cancer and respiratory disease are not the only health risks associated with smoking. More men and women in the United States have died from cardiovascular disease attributed to smoking than cancer.<sup>15</sup> Twenty-one percent of all coronary heart disease deaths in the United States each year are attributable to smoking. This risk is strongly dose-related.<sup>16</sup> Smoking triples the risk of dying from heart disease among middle-aged men and women.<sup>17</sup> Cardiovascular smoking deaths are also due to hypertension and stroke. The risk of ischemic stroke is nearly doubled by smoking.<sup>18</sup> Smoking accounted for 18 percent of all stroke deaths.<sup>19</sup> Both active and passive smoking are associated with an increase in the progression of atherosclerosis.<sup>20</sup> More than 128,000 Americans die from smoking related cardiovascular diseases.<sup>21</sup>

**Other Direct Health Harms From Smoking.** Heart disease, cancer and respiratory diseases are just a few of the physical and medical problems associated with smoking. Smoking may reduce fertility and lead to impotence among men.<sup>22</sup> Cigarette smoking increases both the risk and the severity of rheumatoid arthritis.<sup>23</sup> Hearing loss and vision problems, including cataracts, have been linked to smoking.<sup>24</sup> Chronic coughing, increased phlegm, emphysema and bronchitis have been well-established products of smoking for decades; and smokers are also more susceptible to influenza and more likely to experience severe symptoms when they get the flu.<sup>25</sup> While many smokers believe that smoking relieves stress, it is actually a major cause. Smoking only appears to reduce stress because it lessens the irritability and tension caused by the underlying nicotine addiction.<sup>26</sup>

**Harm Caused by Smokeless Tobacco Use.** The Surgeon General has determined that the use of oral snuff can lead to oral cancer, gum disease, and nicotine addiction.<sup>27</sup> Constant exposure to tobacco juice causes cancer of the esophagus, pharynx, larynx, stomach and pancreas. Smokeless tobacco users are at a heightened risk for oral cancer compared to non-users and these cancers can form within five years of regular use.<sup>28</sup> New studies have found that the levels of the carcinogenic NNK in smokeless tobacco products were comparable to those cigarettes<sup>29</sup> and using Swedish snus can heighten one's risk for pancreatic cancer.<sup>30</sup> A 2008 study from the WHO International Agency for Research on Cancer concluded that smokeless tobacco users have an 80 percent higher risk of developing oral cancer and a 60 percent higher risk of developing pancreatic and esophageal cancer.<sup>31</sup> Spit tobacco causes leukoplakia, a disease of the mouth characterized by white patches and oral lesions on the cheeks, gums, and/or tongue. Leukoplakia, which can lead to oral cancer, occurs in more than half of all users in the first three years of use. Studies have found that 60 to 78 percent of smokeless tobacco users have oral lesions.<sup>32</sup> Gum disease (gingivitis) is also caused by spit tobacco.<sup>33</sup> Spit tobacco has also been linked to dental caries. A study by the National Institutes of Health and the Centers for Disease Control and Prevention found chewing tobacco users were four times more likely than non-users to have decayed dental root surfaces.<sup>34</sup>

**Harms from Pregnant Smokers or Exposure to Secondhand Smoke.** Even more disturbing is the impact of smoking on pregnant women. Research studies have found that smoking and exposure to secondhand smoke among pregnant women is a major cause of spontaneous abortions, stillbirths, and sudden infant death syndrome (SIDS) after birth.<sup>35</sup> According to a meta-analysis of published studies, tobacco use is responsible each year for 19,000 to 141,000 spontaneous abortions, 1,900 to 4,800 infant deaths caused by prenatal or pre-birth disorders, and 1,200 to 2,200 deaths from SIDS.<sup>36</sup> A more recent comprehensive study found that parental smoking causes 2,800 deaths at birth and 2,000 deaths from SIDS.<sup>37</sup> Children exposed to secondhand smoke before and after birth are at a great risk of abnormal blood pressure, cleft palates and lips, childhood leukemia, attention deficit disorder, childhood wheezing and respiratory disorders.<sup>38</sup>

**Other Secondhand Smoke Harms.** Secondhand smoke is the combination of "mainstream smoke" (the smoke exhaled by a smoker) and side-stream smoke (from the burning end of the cigarette). Secondhand smoke is also referred to as environmental tobacco smoke, passive smoke, or involuntary tobacco smoke. It is a complex mixture of over 4,000 chemicals that are produced by the burning materials of a cigarette. Secondhand smoke exposure is causally associated with several different health risks in both children and adults. Children exposed to secondhand smoke are at a higher risk of Sudden Infant Death Syndrome (SIDS), acute lower respiratory tract infections, asthma induction and exacerbation, chronic respiratory symptoms, middle ear infections.<sup>39</sup> In adults, secondhand smoke exposure increases the risk of lung cancer, nasal sinus cancer, heart disease mortality, acute and chronic coronary heart disease morbidity eye and nasal irritation.<sup>40</sup> A 1997 analysis of 37 epidemiological studies of lung cancer and ETS found that lifelong nonsmokers living with smokers had, on average, a 24 percent higher chance of contracting lung cancer than those living with nonsmokers, and that those exposed to the heaviest smokers for the longest time had the highest risks.<sup>41</sup> Subsequent research studies have made similar findings.<sup>42</sup> Secondhand smoke is listed as a carcinogen in the U.S. Public Health Services' *Ninth Report on Carcinogens*, as recommended by a scientific advisory panel of the National Toxicology Program that unanimously affirmed the findings of two other scientific groups that secondhand smoke is a carcinogen and should be included in the report.<sup>43</sup>

**Immediate and Short-Term Harms from Smoking.** While most of the major health harms from smoking, such as lung cancer and heart disease, typically appear after years of tobacco use, many health problems can appear almost immediately, even among otherwise young and healthy kids. For example, cigarette smoking immediately increases heart rate and blood pressure, and the resting heart rates of young adult smokers are two to three beats per minute faster than nonsmokers.<sup>44</sup> In addition, high school seniors who are regular smokers and began smoking by grade nine are 2.4 times more likely than their nonsmoking peers to report poorer overall health. High school seniors who smoke are 2.4 to 2.7 times more likely to report cough with phlegm or blood, shortness of breath when not exercising, and wheezing or gasping.<sup>45</sup> Teens who smoke are also three times more likely than nonsmokers to use alcohol, eight times more likely to use marijuana, and 22 times more likely to use cocaine.<sup>46</sup>

**Tobacco Use and Appearance.** Concern about body weight and appearance are just a few reasons that smoke smokers begin. Most adolescents believe that smoking controls body weight and many times women report that they smoke to keep their weight down.<sup>47</sup> While smoking cessation has been shown to result in weight gain among both men and women, initiation of smoking does not appear to be associated with weight loss.<sup>48</sup> Among women, the average weight of current smokers is only modestly lower than that of never or long-term former smokers. Smoking has also been linked with facial wrinkling. Smokers were significantly more likely than nonsmokers to be evaluated with having prominent wrinkling.<sup>49</sup>

**Smoking Addiction Starts Early.** The peak years for first trying to smoke appear to be in the sixth and seventh grades, or between the ages of 11 and 12, with a considerable number starting even earlier.<sup>50</sup> Within weeks or just days of first starting to smoke occasionally, young smokers show numerous signs of addiction, such as feeling anxious or irritable and having strong urges to smoke;<sup>51</sup> and more than a third of all kids who ever try smoking a cigarette will become regular, daily smokers before they even leave high school.<sup>52</sup> Addiction rates for experimenters who become habitual users for smoking are higher than addiction rates for marijuana, alcohol, or even cocaine.<sup>53</sup> Every day more than 3,500 kids under 18 try smoking for the first time, and another 1,000 kids who have already experimented with cigarettes become new regular daily smokers.<sup>54</sup> Overall, more than 80 percent of all adult smokers first become regular smokers before the age of 18 and more than 90 percent do so before leaving their teens.<sup>55</sup>

**Quitting Is Difficult, But Not Impossible.** Although half of all Americans who have ever smoked have quit, and most current smokers want to stop, an established addiction to nicotine is difficult to escape. Of the more than one million smokers who quit each year, 75 to 80 percent relapse within six months.<sup>56</sup> To quit, smokers must not only overcome their physiological dependence on nicotine but also cut their strong psychological and social ties to smoking or otherwise using tobacco. The three most effective components of smoking cessation treatment are pharmacological treatments (such as nicotine gum and patches), clinician-provided social support and advice, and skills training regarding techniques to achieve and maintain abstinence.<sup>57</sup> Another treatment approach combines nicotine replace, counseling and the use of anti depressants like bupropion. In general, more inclusive treatments are more effective in producing long-term abstinence from tobacco, and combined therapies raise the absolute percentage of smokers who remain abstinent.

**Health Benefits From Quitting.** There are substantial and immediate health benefits from quitting smoking. A 2007 study in the *New England Journal of Medicine* found that 11.7 percent of the decrease in coronary heart disease deaths between 1980 and 2000 were avoided or postponed by quitting smoking.<sup>58</sup> Upon quitting, former smokers' blood circulation immediately increases, their blood pressure and heart rate quickly return to normal, and the carbon monoxide and oxygen levels in the blood soon return to normal. Within a few days of quitting, a person's breathing becomes easier and their sense of smell and taste improve.<sup>59</sup> One year after quitting, a person's additional risk of heart disease is reduced by half, and after 15 years, this risk equals that of a person who never smoked.<sup>60</sup> Five to 15 years after quitting, the risk of stroke for an ex-smoker equals that of a person who never smoked.<sup>61</sup> Within 10 years of quitting a former smokers risk of developing lung cancer is 30 to 50 percent below that of a person who continues to smoke. The risk of developing cancers of the mouth, throat and esophagus lessen significantly after five years of quitting.<sup>62</sup> Persons aged 60 to 64 years of age who quit smoking are 10 percent less likely to die during the next 15 years than regular smokers are.<sup>63</sup> The benefit is even greater for individuals who quit smoking before the age of 50. Their risk of dying in the next 15 years is half that of a person who smokes.<sup>64</sup>

Despite the numerous reports of the Surgeon General and the National Institute for Health on the risks of smoking, more than 45 million Americans still smoke. Approximately half of all continuing, long-term smokers will die prematurely as a result of their addiction. These tobacco-related deaths and illnesses are preventable.

**Campaign for Tobacco-Free Kids, March 23, 2011 / Ann Boonn**

More information on the health harms of tobacco use is available at  
[http://www.tobaccofreekids.org/facts\\_issues/fact\\_sheets/toll/products/](http://www.tobaccofreekids.org/facts_issues/fact_sheets/toll/products/).

- <sup>1</sup> U.S. Centers for Disease Control and Prevention (CDC), "Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses – United States, 2000-2004," *Morbidity and Mortality Weekly Report (MMWR)* 57(45), November 14, 2008 <http://www.cdc.gov/mmwr/PDF/wk/mm5745.pdf>. (AIDS) CDC, "Table 7. Estimated numbers of deaths of persons with AIDS, by year of death and selected characteristics, 2001–2005 and cumulative—United States and dependent areas," HIV/AIDS Surveillance Report, Volume 17, Revised Edition, June 2007, <http://www.cdc.gov/hiv/topics/surveillance/resources/reports/2005report/pdf/table7.pdf>; (Alcohol) Mokdad, AH, et al., "Actual Causes of Death in the United States, 2000," *Journal of the American Medical Association (JAMA)* 291(10):1238-1245, March 10, 2004 [with correction in *JAMA* 293(3):298, January 19, 2005]; (Motor vehicle) National Highway Traffic Safety Administration's National Center for Statistics and Analysis, *2006 Traffic Safety Annual Assessment – A Preview*, DOT HS 810 791, July 2007, <http://www-nrd.nhtsa.dot.gov/pubs/810791.pdf>; (Homicide, Suicide, Drug-Induced) Kung HC, et al., "Preliminary Data for 2005," *Health E-Stats*, National Center for Health Statistics, September 2007, [http://www.cdc.gov/nchs/data/hestat/prelimdeaths05/preliminarydeaths05\\_tables.pdf#2](http://www.cdc.gov/nchs/data/hestat/prelimdeaths05/preliminarydeaths05_tables.pdf#2).
- <sup>2</sup> National Cancer Institute (NCI), *Changes in Cigarette-Related Disease Risks and Their Implications for Prevention and Control*, Monograph 8, National Institutes of Health (NIH), 1997.
- <sup>3</sup> Available at <http://profiles.nlm.nih.gov/NN/B/C/X/B/>.
- <sup>4</sup> CDC, "Vital Signs: Current Cigarette Smoking Among Adults Aged ≥ 18 Years - United States, 2009," *MMWR* 59(35):1135-1140, September 7, 2010. <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5935a3.htm>.
- <sup>5</sup> Tverdal, A & Bjartveit, K, "Health consequences of pipe versus cigarette smoking," *Tobacco Control*, published online October 15, 2010. NCI, *Cigars: Health Effects and Trends*, Smoking and Tobacco Control Monograph No. 9, 1998, <http://cancercontrol.cancer.gov/tcrb/monographs/9/index.html>. See also, Campaign for Tobacco-Free Kids, *The Rise of Cigars and Cigar-Smoking Harms*, <http://www.tobaccofreekids.org/research/factsheets/pdf/0333.pdf>.
- <sup>6</sup> U.S. Department of Health and Human Services (HHS), *The Health Consequences of Smoking. A Report of the Surgeon General*, 2004, <http://www.surgeongeneral.gov/library/smokingconsequences/>. See also, Thun, M, "Mixed progress against lung cancer," *Tobacco Control* 7:223-226, 1998.
- <sup>7</sup> CDC, "Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Economic Costs—United States 1995-1999," *MMWR*, April 11, 2002, [www.cdc.gov/mmwr/preview/mmwrhtml/mm5114a2.htm](http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5114a2.htm).
- <sup>8</sup> HHS, *The Health Consequences of Smoking. A Report of the Surgeon General*, 2004, <http://www.surgeongeneral.gov/library/smokingconsequences/>.
- <sup>9</sup> HHS, *Reducing the Health Consequences of Smoking: 25 Years of Progress. A Report of the Surgeon General*, HHS Publication No 89-8911, 1989, <http://profiles.nlm.nih.gov/NN/B/B/X/S/>.
- <sup>10</sup> HHS, *The Health Consequences of Smoking. A Report of the Surgeon General*, 2004.
- <sup>11</sup> CDC, "Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses – United States, 2000-2004," *MMWR* 57(45), November 14, 2008.
- <sup>12</sup> American Lung Association Smoking Fact Sheet, September 2000.
- <sup>13</sup> HHS, *The Health Consequences of Smoking. A Report of the Surgeon General*, 2004.
- <sup>14</sup> HHS, *The Health Consequences of Smoking. A Report of the Surgeon General*, 2004. See also, CDC, "Cigarette smoking-attributable mortality and years of potential life lost—United States, 1990," *MMWR* 42(33):645-8, 1993, <http://www.cdc.gov/mmwr/PDF/wk/mm4233.pdf>.
- <sup>15</sup> CDC, "Cigarette smoking-attributable mortality and years of potential life lost—United States, 1990," *MMWR* 42(33):645-8, 1993, <http://www.cdc.gov/mmwr/PDF/wk/mm4233.pdf>.
- <sup>16</sup> HHS, *Reducing the Health Consequences of Smoking: 25 Years of Progress. A Report of the Surgeon General*, 1989, <http://profiles.nlm.nih.gov/NN/B/B/X/S/>.
- <sup>17</sup> CDC, "Cigarette smoking-attributable mortality and years of potential life lost—United States, 1990," *MMWR* 42(33):645-8, 1993, <http://www.cdc.gov/mmwr/PDF/wk/mm4233.pdf>.
- <sup>18</sup> Shinton, R & Bevers, G, "Meta-analysis of relation between smoking and stroke," *BMJ* 298:789-794, 1989.
- <sup>19</sup> HHS, *Reducing the Health Consequences of Smoking: 25 Years of Progress. A Report of the Surgeon General*, 1989, <http://profiles.nlm.nih.gov/NN/B/B/X/S/>.
- <sup>20</sup> Howard, G, et al., "Cigarette smoking and progression of atherosclerosis," *JAMA* 279:119-124, 1998.
- <sup>21</sup> CDC, "Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses – United States, 2000-2004," *MMWR* 57(45), November 14, 2008.
- <sup>22</sup> On early menopause, see, e.g., Cramer, DW, et al., "Cross-sectional and Case-controlled Analyses of the Association Between Smoking and Early Menopause," *Maturitas* 22(2):79-87, September 1995. On incontinence, see, e.g., Tampakoudis, P, et al., "Cigarette Smoking and Urinary Incontinence in Women – A New Calculative Method of Estimating the Exposure to Smoke," *European Journal of Gynecology and Reproductive Biology* 63(1):27-30, November 1995. On reduced female fertility, see, e.g., Van Voorhis, BJ, et al., "The Effects of Smoking on Ovarian Function and Fertility During Assisted Reproduction Cycles," *Obstetric Gynecology* 88(5):785-91, November 1996. On reduced male infertility see, e.g., Close, CE, et al., "Cigarettes, Alcohol, and Marijuana are Related to Pyospermia in Infertile Men," *Journal of Urology* 144(4):900-03, October 1990; Moskova, P & Popov, I, "Sperm Quality in Smokers and Nonsmokers Among Infertile Families," *Akusherstvo I Ginekologija* 32(1):28-30, 1993 [in Bulgarian].
- <sup>23</sup> Saag, KG, et al., "Cigarette Smoking and Rheumatoid Arthritis Severity," *Annals of Rheumatic Diseases* 56(8):463-69, August 1997; Silman, AJ, et al., "Cigarette Smoking Increases the Risk of Rheumatoid Arthritis: Results from a Nationwide Study of Disease-Discordant Twins," *Arthritis and Rheumatism* 39(5):732-35, May, 1996.
- <sup>24</sup> Noorhassim, I & Rampal, KG, "Multiplicative Effect of Smoking and Age on Hearing Impairment," *American Journal of Otolaryngology* 19(4):240-43, July-August 1998; Cruickshanks, KJ, et al., "Cigarette Smoking and Hearing Loss: The Epidemiology of Hearing Loss Study,"



- JAMA 279(21):1715-19, June 3, 1998. For eye problems, see, e.g., Klein, R, et al., "Relation of Smoking to the Incidence of Age-Related Maculopathy: The Beaver Dam Eye Study," *American Journal of Epidemiology (AJE)* 147(2):103-10, January 15, 1998; Cumming, RG & Mitchell, P, "Alcohol, Smoking, and Cataracts," *Archives of Ophthalmology* 115(10):1296-303, October 1997; Sibony, PA, et al., "The Effects of Tobacco Smoking on Smooth Pursuit Eye Movements," *Annals of Neurology* 23(3):238-41, March 1988;
- <sup>25</sup> Kark, JD, et al., "Cigarette smoking as a factor of epidemic influenza in young men," *New England Journal of Medicine (NEJM)* 307(17):1042-46, October 21, 1982; JD, Kark & Lebiush, M, "Smoking and epidemic influenza like illness in female military recruits: a brief survey," *American Journal of Public Health (AJPH)* 71(5):530-32, May 1981.
- <sup>26</sup> Parrott, AC, "Does Cigarette Smoking Cause Stress," *American Psychologist* 54(10):817-20, October 1999.
- <sup>27</sup> HHS, *The Health Consequences of Using Smokeless Tobacco: A Report of the Advisory Committee to the Surgeon General*, April 1986, <http://profiles.nlm.nih.gov/NN/B/B/F/C/>.
- <sup>28</sup> The S.T.O.P. Guide (The Smokeless Tobacco Outreach and Prevention Guide): A Comprehensive Directory of Smokeless Tobacco Prevention and Cessation Resources, Applied Behavioral Science Press, 1997; Hatsukami, D & Severson, H, "Oral Spit Tobacco: Addiction, Prevention and Treatment." *Nicotine and Tobacco Research*, 1:21-44, 1999.
- <sup>29</sup> Hecht, SS, et al., "Similar Exposure to a Tobacco-Specific Carcinogen in Smokeless Tobacco Users and Cigarette Smokers," *Cancer Epidemiology Biomarkers & Prevention* 16(8):1-6, 2007.
- <sup>30</sup> Luo, J, et al., "Oral use of Swedish moist snuff (snus) and risk for cancer of the mouth, lung, and pancreas in male construction workers: a retrospective cohort study," *The Lancet*, May 10, 2007.
- <sup>31</sup> Boffetta, P, et al., "Smokeless tobacco and cancer," *The Lancet* 9:667-675, 2008.
- <sup>32</sup> The S.T.O.P. Guide (The Smokeless Tobacco Outreach and Prevention Guide): A Comprehensive Directory of Smokeless Tobacco Prevention and Cessation Resources, Applied Behavioral Science Press, 1997.
- <sup>33</sup> The S.T.O.P. Guide (The Smokeless Tobacco Outreach and Prevention Guide): A Comprehensive Directory of Smokeless Tobacco Prevention and Cessation Resources, Applied Behavioral Science Press, 1997.
- <sup>34</sup> Tomar, SL, "Chewing Tobacco Use and Dental Caries Among U.S. Men," *Journal of the American Dental Association* 130:160, 1999.
- <sup>35</sup> On spontaneous abortions, see, e.g., Mendola, P, et al., "Risk of Recurrent Spontaneous Abortion, Cigarette Smoking, and Genetic Polymorphisms in NAT2 and GSTM1," *Epidemiology* 9(6):666-668, November 1999; Shiverick, KT & Salafia, C, "Cigarette Smoking and Pregnancy I: Ovarian, Uterine and Placental Effects," *Placenta* 20(4):265-272, May 1999; Ness, RB, et al., "Cocaine and Tobacco Use and the Risk of Spontaneous Abortion," *NEJM* 340(5):333-339, February 4, 1999; Chatenoud, L, et al., "Paternal and Maternal Smoking Habits Before Conception and During the First Trimester: Relation to Spontaneous Abortions," *Annals of Epidemiology* 8(8):520-26, November 1998; Hrubá, D & Kachlik, P, "Relation Between Smoking in Reproductive-Age Women and Disorders in Reproduction," *Ceska Gynekol* 62(4):191-196, August 1997 [in Czech]; Domínguez-Rojas, V, et al., "Spontaneous Abortion in a Hospital Population: Are Tobacco and Coffee Intake Risk Factors?," *European Journal of Epidemiology* 10(6):665-668, December 1994; Walsh, RA, "Effects of Maternal Smoking on Adverse Pregnancy Outcomes: Examination of the Criteria for Causation," *Human Biology* 66(6):1059-1092, December 1994; Windham, GC, et al., "Parental Cigarette Smoking and the Risk of Spontaneous Abortion," *AJE* 135(12):1394-403, June 15, 1992; Armstrong, BG, et al., "Cigarette, Alcohol, and Coffee Consumption and Spontaneous Abortion," *AJPH* 82(1):85-87, January 1992; Pattinson, HA, et al., "The Effect of Cigarette Smoking on Ovarian Function and Early Pregnancy Outcome Of In Vitro Fertilization Treatment," *Fertility and Sterility* 55(4):780-783, April 1991; Ahlberg, G, Jr. & Bodin, L, "Tobacco Smoke Exposure and Pregnancy Outcome Among Working Women: A Prospective Study At Prenatal Care Centers In Orebro County, Sweden," *AJE* 133(4):338-347, February 1991; Himmelberger, DU, et al., "Cigarette Smoking During Pregnancy and the Occurrence of Spontaneous Abortion and Congenital Abnormality," *AJE* 108(6):470-479, December 1978; Kline, J, et al., "Smoking: A Risk Factor for Spontaneous Abortions," *NEJM* 291(15):793-96, October 1977. See also, Kline, J, et al., "Cigarette Smoking and Spontaneous Abortion of Known Karyotype: Precise Data But Uncertain Inferences," *AJE* 141(5):417-427, March 1995; Economides, D & Braithwaite, J, "Smoking, Pregnancy, and the Fetus," *Journal of the Royal Society of Health* 114(4):198-201, August 1994; Fredricsson, B & Gilljam, H, "Smoking and Reproduction: Short and Long Term Effects and Benefits of Smoking Cessation," *Acta Obstetrica Gynecologica Scandinavica* 71(8):580-592, December 1992. But see, also, Windham, GC, et al., "Exposure to Environmental and Mainstream Tobacco Smoke and Risk of Spontaneous Abortion," *AJE* 149(3):243-247, February 1, 1999; Sandahl, B, "Smoking Habits and Spontaneous Abortion," *European Journal of Obstetric Gynecology and Reproductive Biology* 31(1):23-31, April 1989. On still births, see, e.g., Raymond, EG, et al., "Effects of Maternal Age, Parity, and Smoking on the Risk of Stillbirth," *British Journal of Obstetric Gynaecology* 101(4):301-306, April 1994; Ahlberg, G, Jr. & Bodin, L, "Tobacco Smoke Exposure and Pregnancy Outcome Among Working Women: A Prospective Study At Prenatal Care Centers In Orebro County, Sweden," *AJE* 133(4):338-347, February 1991. On sudden infant death syndrome, see, e.g., Cooke, RW, "Smoking, Intra-Uterine Growth Retardation and Sudden Infant Death Syndrome," *International Journal of Epidemiology* 27(2):238-41, April 1998.
- <sup>36</sup> DiFranza, JR & Lew, RA, "Effect of Maternal Cigarette Smoking on Pregnancy Complications and Sudden Infant Death Syndrome," *Journal of Family Practice* 40(4):385-94, April 1995.
- <sup>37</sup> Aligne, CA & Stoddard, JJ, "Tobacco and Children: An Economic Evaluation of the Medical Effects of Parental Smoking," *Archives of Pediatric and Adolescent Medicine* 151(7):648-53, July 1997.
- <sup>38</sup> On abnormal blood pressure in infants and children, see, e.g., Morley, R, et al., "Maternal Smoking and Blood Pressure in 7.5 to 8 Year Old Offspring," *Archives of Disease in Childhood* 72(2):120-24, February 1995; Blake, KV, et al., "Maternal Cigarette Smoking During Pregnancy, Low Birth Weight and Subsequent Blood Pressure in Early Childhood," *Early Human Development* 57:137-147, 2000. On cleft palates and lips, see Nagourney, E, "Consequences: Linking Cleft Palates and Smoking Moms," *New York Times*, April 12, 2000 [citing recent study in *Plastic and Reconstructive Surgery*]. On links with carcinogens and cancer, see, e.g., Lackmann, GM, et al., "Metabolites of a Tobacco-Specific Carcinogen in Urine from Newborns," *Journal of the National Cancer Institute (JNCI)* 91(5):459-65, March 1999; Finette, BA, et al., "Gene Mutations with Characteristic Deletions in Cord Blood T Lymphocytes Associated with Passive Maternal Exposure to Tobacco Smoke," *Nature Medicine* 4(10):1144-51, October 1998; Wise, J, "Carcinogen in Tobacco Smoke Can Be Passed to Fetus," *British Medical Journal (BMJ)*

- 317(7158):555, August 29, 1998; Crawford, FG, et al., "Biomarkers of Environmental Tobacco Smoke in Preschool Children and Their Mothers," *JNCI* 86(18):1398-402, September 21, 1994; Stjernfeldt, M, et al., "Maternal Smoking and Irradiation During Pregnancy as Risk Factors for Child Leukemia," *Cancer Detection and Prevention* 16(2):129-35, 1992. On birth defects, see, e.g., Kallen, K, "Maternal Smoking During Pregnancy and Limb Reduction Malformations in Sweden," *AJPH* 87(1):29-32, January 1997; Czeizel, AE, et al., "Smoking During Pregnancy and Congenital Limb Deficiency," *BMJ* 308(6942):1473-76, June 1994; Drews, CD, et al., "The Relationship Between Idiopathic Mental Retardation and Maternal Smoking During Pregnancy," *Pediatrics* 97(4):547-53, April 1997. On colic, see Reijneveld, SA, et al., "Infantile Colic: Maternal Smoking As Potential Risk Factor," *Archives of Disease in Childhood* 83:302-303, October 2000. On wheezing and respiratory problems, see, e.g., Hu, FB, et al., "Prevalence of Asthma and Wheezing in Public Schoolchildren: Association with Maternal Smoking During Pregnancy," *Annals of Allergy, Asthma, and Immunology* 79(1):80-84, July 1997; Tager, IB, et al., "Maternal Smoking During Pregnancy: Effects on Lung Function During the First 18 Months of Life," *American Journal of Respiratory and Critical Care Medicine* 152(3):977-83, September 1995; Lux, AL, et al., "Wheeze Associated with Prenatal Tobacco Smoke Exposure: A Prospective, Longitudinal Study," *Archives of Disease in Childhood* 83:307-12, October 2000.
- <sup>39</sup> California Environmental Protection Agency (CEPA), *Health Effects of Exposure to Environmental Tobacco Smoke*, NIH, 1999.
- <sup>40</sup> CEPA, *Health Effects of Exposure to Environmental Tobacco Smoke*, NIH, 1999.
- <sup>41</sup> Hackshaw, AK, Law, MR, & Wald, NJ, "The Accumulated Evidence on Lung Cancer and Environmental Tobacco Smoke," *BMJ* 315:980-988, October 18, 1997.
- <sup>42</sup> Boffetta, P, et al., "Multicenter Case-Control Study of Exposure to Environmental Tobacco Smoke and Lung Cancer in Europe," *JNCI* 90:1440-50, October 7, 1998. See also, CEPA, *Health Effects of Exposure to Environmental Tobacco Smoke*, 1997.
- <sup>43</sup> National Toxicology Program, Public Health Service, *Ninth Report on Carcinogens*, 2000, <http://ehis.niehs.nih.gov/roc/toc9.html#toc>; Robertson, GD, "Panel Says Secondhand Smoke, Alcohol Should Be Considered Carcinogens," *Associated Press*, December 3, 1998.
- <sup>44</sup> Royal College of Physicians of London, "Smoking and the young: a report of a working party of the Royal College of Physicians," Oxford: 33, 1992; HHS, *Preventing Tobacco Use Among Young People: A Report of the Surgeon General*, Washington, DC: HHS, Public Health Service, Office of the Surgeon General, 1994, [http://www.cdc.gov/tobacco/data\\_statistics/sgr/sgr\\_1994/index.htm](http://www.cdc.gov/tobacco/data_statistics/sgr/sgr_1994/index.htm).
- <sup>45</sup> Arday DR, et al., "Cigarette smoking and self-reported health problems among US high school seniors, 1982-1989," *American Journal of Health Promotion*, 10(2):111-116, 1995.
- <sup>46</sup> HHS, *Preventing Tobacco Use Among Young People: A Report of the Surgeon General*, 1994, [http://www.cdc.gov/tobacco/data\\_statistics/sgr/sgr\\_1994/index.htm](http://www.cdc.gov/tobacco/data_statistics/sgr/sgr_1994/index.htm).
- <sup>47</sup> HHS, *Women and Smoking: A Report of the Surgeon General*, 2001, [http://www.cdc.gov/tobacco/data\\_statistics/sgr/sgr\\_2001/index.htm](http://www.cdc.gov/tobacco/data_statistics/sgr/sgr_2001/index.htm).
- <sup>48</sup> HHS, *Women and Smoking: A Report of the Surgeon General*, 2001, [http://www.cdc.gov/tobacco/data\\_statistics/sgr/sgr\\_2001/index.htm](http://www.cdc.gov/tobacco/data_statistics/sgr/sgr_2001/index.htm).
- <sup>49</sup> HHS, *Women and Smoking: A Report of the Surgeon General*, 2001, [http://www.cdc.gov/tobacco/data\\_statistics/sgr/sgr\\_2001/index.htm](http://www.cdc.gov/tobacco/data_statistics/sgr/sgr_2001/index.htm).
- <sup>50</sup> NIH, *National Survey Results on Drug Use from the Monitoring the Future Study, 1975-1997*, 1998.
- <sup>51</sup> DiFranza, JR, et al., "Initial Symptoms of Nicotine Dependence in Adolescents," *Tobacco Control* 9:313-9, September 2000.
- <sup>52</sup> CDC, "Selected Cigarette Smoking Initiation and Quitting Behaviors Among High School Students—United States - 1997," *MMWR* 47(19):386-389, May 22, 1998, <http://www.cdc.gov/mmwr/preview/mmwrhtml/00052816.htm>.
- <sup>53</sup> CDC, "Symptoms or Substance Dependence Associated with Use of Cigarettes, Alcohol, and Illicit Drugs—United States 1991-1992," *MMWR* 44(44):830-931, November 10, 1995, <http://www.cdc.gov/mmwr/preview/mmwrhtml/00039501.htm>; See also the National Survey on Drug Use and Health (NSDUH) and the Monitoring the Future Study for addiction information.
- <sup>54</sup> Substance Abuse and Mental Health Services Administration, (SAMHSA), HHS, *Results from the 2007 National Survey on Drug Use and Health, NSDUH: Detailed Tables*, <http://www.oas.samhsa.gov/NSDUH/2k7NSDUH/tabs/Sect4peTabs10to11.pdf>.
- <sup>55</sup> SAMHSA, HHS, Calculated based on data in 2006 National Household Survey on Drug Use and Health.
- <sup>56</sup> Carmody, TP, "Preventing relapse in the treatment of nicotine addiction: Current issues and future directions," *Journal of Psychoactive Drugs* 24(2):131-58, 1992.
- <sup>57</sup> National Institute on Drug Abuse, NIH, *Addicted to Nicotine: A National Research Forum*, July 27-28, 1998.
- <sup>58</sup> Ford, ES, et al., "Explaining the decrease in U.S. deaths from Coronary Disease, 1980-2000," *NEJM* 356(23):2388-2398, June 7, 2007.
- <sup>59</sup> HHS, *The Health Benefits of Smoking Cessation: A Report of the Surgeon General*, HHS Publication No. (CDC) 90-8416, 1990, <http://profiles.nlm.nih.gov/NN/B/B/C/T/>.
- <sup>60</sup> HHS, *Reducing the Health Consequences of Smoking: 25 Years of Progress. A Report of the Surgeon General*, 1989, <http://profiles.nlm.nih.gov/NN/B/B/X/S/>.
- <sup>61</sup> HHS, *Reducing the Health Consequences of Smoking: 25 Years of Progress. A Report of the Surgeon General*, 1989, <http://profiles.nlm.nih.gov/NN/B/B/X/S/>.
- <sup>62</sup> HHS, *The Health Benefits of Smoking Cessation: A Report of the Surgeon General*, HHS Publication No. (CDC) 90-8416, 1990, <http://profiles.nlm.nih.gov/NN/B/B/C/T/>.
- <sup>63</sup> HHS, *The Health Benefits of Smoking Cessation: A Report of the Surgeon General*, HHS Publication No. (CDC) 90-8416, 1990, <http://profiles.nlm.nih.gov/NN/B/B/C/T/>.
- <sup>64</sup> HHS, *The Health Benefits of Smoking Cessation: A Report of the Surgeon General*, HHS Publication No. (CDC) 90-8416, 1990, <http://profiles.nlm.nih.gov/NN/B/B/C/T/>.