

Study Design and Analytic Epidemiology for Outbreak Investigations

Rosalie Trevejo, DVM, MPVM, PhD
Oregon Public Health Division
Acute and Communicable Disease Prevention



1

Objectives

- Define analytic epidemiology
- Describe types of studies to investigate outbreaks
- Discuss control group selection
- Prepare 2 x 2 table
- Calculate and interpret measures of association




2

2

Analytic Epidemiology

- Differs from descriptive epi (person, place, time)
- Develop then test hypothesis
- Examine **exposures** and **outcomes**
- Use **comparison group** (controls)



3

3

Exposure and Outcome

- Exposure= potential risk factor
 - Food item
 - Occupational
 - Behavioral (e.g., smoking)
- Outcome= Disease
 - Cancer
 - Diarrhea
 - Injury

Oregon
Health
Authority

4

4

Common Types of Studies

- Cohort
- Case-Control
- Case-case
- These are **observational** studies, as compared with experimental studies

Cohort Design

- Start with defined group, classified by exposure status
 - Exposed group
 - Non-exposed group
- Assess who developed disease

Cohort Studies examples

- Nurse's Health Study- 1970's
120,000 female nurses – risk factors for cancer and cardiovascular disease
- Outbreak following a wedding reception
- Illness in a school classroom
- Outbreak after church potluck

www.nhs3.org

Oregon
Health
Authority

7

7

Why use a cohort?

- Known denominator: # at **risk** of disease
- Compare exposed and unexposed groups
- However, not always possible to define cohort




Oregon
Health
Authority


8

8

Cohort Study



Cohort Study



Case-Control Design

- Classify people by outcome
 - Cases: have disease
 - Controls: do not have disease
- Assess past exposure

Case-control study examples

- Smoking and lung cancer, 1950s
- *Salmonella* outbreak associated with nationally distributed product
- Restaurant outbreak of norovirus




Selecting Controls

- Do not have the disease
- At risk of disease: had potential for exposure
- Should be representative of cases
- Key: **If they developed the disease, could they be identified as case?**

Potential control groups


- Others who ate at same restaurant
- Neighborhood residents
- Family members
- Friends
- Coworkers
- Random digit dialing
- Classmates
- Medical clinic patients

Case-control Study



15

Case-control Study Example



16

Advantages & Disadvantages

	Cohort	Case-control
Advantages	Can evaluate rare exposures Can calculate risk	Quick Smaller size Cost effective Can evaluate rare disease
Disadvantages	Inefficient if incubation is long Potential expense Loss to follow up	Recall bias Selection bias

Measures of Association

- Quantifies magnitude of association between exposure and disease
 - Cohort study: relative risk (RR)
 - Case-control study: odds ratio (OR)

2 X 2 table

	Diseased	Not diseased	Row total
Exposed	a	b	a+b
Not exposed	c	d	c+d
Column total	a+c	b+d	a+b+c+d= Grand total

Cohort Study: Relative Risk (RR)

- Risk of disease in exposed $\div$ risk of disease in unexposed
- $RR = a/(a+b) \div c/(c+d)$

		Disease	
		Yes	No
Exposure	Yes	a	b
	No	c	d

How to interpret If $RR = 1$?

Calculate RR for Cohort Study

- *Salmonella* Enteritidis outbreak among group of coworkers that had catered lunch with taco bar
- 40 workers ate fresh salsa; 30 developed *Salmonella* Enteritidis and 10 did not
- Another 40 workers did not eat fresh salsa; 2 developed *Salmonella* Enteritidis and 38 did not

Relative Risk 2 X 2 table

	<i>Salmonella</i> Enteritidis	No <i>Salmonella</i> Enteritidis	Total
Salsa	a	b	a+b
No salsa	c	d	c+d

Fill in table and calculate RR
Put your answer in chat

2X2 table results

	<i>Salmonella</i> Enteritidis	No <i>Salmonella</i> Enteritidis	Total
Salsa	30	10	40
No salsa	2	38	40

$$RR = a/(a+b) \div c/(c+d)$$

$$30/(30+10) \div 2/(2+38)$$

$$RR = 15$$

Relative Risk conclusion:

Those who ate salsa had higher risk (15X) of developing *Salmonella* Enteritidis than those who didn't eat salsa



Case-Control Study: Odds Ratio (OR)

- Compare odds of exposure for cases and controls

- **Odds Ratio (OR) =**

- $a/c \div b/d$

- ad/bc

		Disease	
		Yes	No
Exposure	Yes	a	b
	No	c	d

Cross product

Oregon
Health
Authority

25

25

Case-Control Study: Calculate OR

- Cases: 40 restaurant patrons with *Salmonella* Enteritidis; 30 ate eggs and 10 did not
- Controls: 40 restaurant patrons without *Salmonella* Enteritidis; 2 ate eggs and 38 did not

Why might a case-control study be good option for restaurant outbreak?

Oregon
Health
Authority

26

26

Odds Ratio 2 X 2 Table

	<i>Salmonella</i> Enteritidis	No <i>Salmonella</i> Enteritidis
Eggs	a	b
No eggs	c	d
Total	a+c	b+d

Fill in table and calculate OR
Put your answer in chat

Oregon
Health
Authority

27

27

Odds ratio 2X2 table results

- Odds Ratio (OR) = $(a*d)/(b*c)$
- OR=
- Conclusion:
Cases were ____ times
more likely than controls to
have eaten eggs



Complete with your table group.

Oregon
Health
Authority

29

29

Case-case study example: Shotgun

- Compare exposures between 2 groups of cases
- Cluster of STEC 4 cases: compare exposure frequencies with background exposure frequencies in Shotgun database
- Shotgun database provides background exposure frequencies from previously interviewed cases (Salmonella and STEC)

Exposure Code	YES	%	Backgrnd	Binomial
331 CarrotsLoose	2 / 4	50	18%	0.1547
336 PepperBell	2 / 4	50	26%	0.2813
338 PepperRed	2 / 4	50	16%	0.1180
339 PepperYellow	2 / 4	50	9%	0.0456

31

health
Authority

31

Is this a true association?

- P-value <0.05)
<https://tidsskriftet.no/en/2015/09/why-p-value-significant>

- Case control study example
OR=57 (95% CI, 12–280) P<0.0001
<https://www.medcalc.org/calc/index.php>



32


Oregon
Health
Authority

32

Study Design and Analytic Epidemiology

Questions?

**Stand up and stretch
break**



Oregon
Health
Authority

33

33

Study Design and Analytic Epidemiology

Exercise

- Handout: CD 303 Exercises (pages 6-8)
- Complete each question before proceeding to next question
- Review questions together at end

Oregon
Health
Authority

34

34

Study Design and Analytic Epidemiology

Review of Opening Scenario

On **August 10**, the Oregon County Health Department received a report of a child with *E. coli* O157 (O157) diarrhea. This was their first O157 report in over a year. Within 24 hours Oregon County had received 3 more reports of O157 infection in children.

All 4 ill children that were reported to Oregon County Health Department over 24-hour period had attended the Oregon County Fair and had bloody diarrhea. The fair had been held **August 1 – 6**. Over ~120,000 people had attended this fair. However, the fair ended 3 days earlier. The fairground facilities are used for activities throughout the year.



35

35

Study Design and Analytic Epidemiology

Additional Information:

The Oregon County Fair is a major annual event in Oregon each August. It is a large agricultural fair with hundreds of animal exhibits, food vendors, and amusement rides. More than 170,000 visits were recorded this year.

The fairgrounds are supplied by shallow wells approximately 20-feet deep. Since the fairgrounds are only in operation 42 days of year, the fairground water supply is not considered a community water system but is considered an alternative water source and is regulated under the food code. This type of well must be tested within 60 days of an event. Testing for coliforms was performed on June 20th and was negative.



36

36

Study Design and Analytic Epidemiology

Question 1: What hypotheses do you want to evaluate?

37

Oregon
Health
Authority

37

Study Design and Analytic Epidemiology

Additional information

An environmental investigation of the fairground revealed that much of the fair was supplied by chlorinated water. However, in one area of the fairground, a shallow well (well #6) provided unchlorinated water to several vendors who used the water to make beverages and ice. This area was close to the large animal barn.

38

Oregon
Health
Authority

38

Study Design and Analytic Epidemiology

Question 2: Do we need a study, if so what type of study?

Why might case control study be used in this investigation?

39

Oregon
Health
Authority

39

Study Design and Analytic Epidemiology

Additional information

Cases eligible for inclusion in case control study were people who had attended the Fair and who had stool culture positive for *E. coli* O157 and were the first case in a household (primary cases).

Why limit to first case in household?

40

Oregon
Health
Authority

40

Study Design and Analytic Epidemiology

Question 3: Whom might you get for controls? How do you find them?

41



41

Study Design and Analytic Epidemiology

Additional information

For the Oregon County Fair, controls were obtained from credit card receipts of attendees and from persons entered in photography exhibit and children's art exhibit. Controls were selected who had a similar age distribution as the cases.

42



42

Study Design and Analytic Epidemiology

Results

Complete 2X2 tables for each exposure.

Table 1: exposure to water from well #6

	Cases (ill)	Controls (not ill)
Yes, exposed		
No, not exposed		
Total		

Table 2: exposure to Vendor A chicken

	Cases (ill)	Controls (not ill)
Yes, exposed		
No, not exposed		
total		

Study Design and Analytic Epidemiology

Question 4: What is an appropriate measure of association for a case-control study? (put in chat)

- Odds ratio
- $\frac{a}{c} = \frac{ad}{bc}$
- $\frac{b}{d} = \frac{ad}{bc}$
- Measure of association for water from well #6? (put in chat)
 - OR=?
- Measure of association for Vendor A chicken? (put in chat)
 - OR=?

Calculating odds ratios $(a*d)/(b*c)$

Exposure to water from well #6

Exposure to Vendor A chicken