

# Study Design and Analytic Epidemiology for Outbreak Investigations

Rosalie Trevejo, DVM, MPVM, PhD  
Oregon Public Health Division  
Acute and Communicable Disease Prevention



1

## Objectives

- Define analytic epidemiology
- Describe types of studies to investigate outbreaks
- Discuss control group selection
- Prepare 2 x 2 table
- Calculate and interpret measures of association




2

2

## Analytic Epidemiology

- Differs from descriptive epi (person, place, time)
- Develop then test hypothesis
- Examine **exposures** and **outcomes**
- Use **comparison group** (controls)



3

3

## Exposure and Outcome

- Exposure= potential risk factor
  - Food item
  - Occupational
  - Behavioral (e.g., smoking)
- Outcome= Disease
  - Cancer
  - Diarrhea
  - Injury

Oregon  
Health  
Authority

4

4

## Common Types of Studies

---

- Cohort
- Case-Control
- Case-case
- These are **observational** studies, as compared with experimental studies

## Cohort Design

---

- Start with defined group, classified by exposure status
  - Exposed group
  - Non-exposed group
- Assess who developed disease

## Cohort Studies examples

- Nurse's Health Study- 1970's  
120,000 female nurses – risk factors for cancer and cardiovascular disease
- Outbreak following a wedding reception
- Illness in a school classroom
- Outbreak after church potluck

[www.nhs3.org](http://www.nhs3.org)

Oregon  
Health  
Authority

7

7

## Why use a cohort?

- Known denominator: # at **risk** of disease
- Compare exposed and unexposed groups
- However, not always possible to define cohort




Oregon  
Health  
Authority


8

8

## Cohort Study



## Cohort Study



## Case-Control Design

- Classify people by outcome
  - Cases: have disease
  - Controls: do not have disease
- Assess past exposure

## Case-control study examples

- Smoking and lung cancer, 1950s
- *Salmonella* outbreak associated with nationally distributed product
- Restaurant outbreak of norovirus



## Selecting Controls

---


- Do not have the disease
- At risk of disease: had potential for exposure
- Should be representative of cases
- Key: **If they developed the disease, could they be identified as case?**

## Potential control groups

---

- Others who ate at same restaurant
- Neighborhood residents
- Family members
- Friends
- Coworkers
- Random digit dialing
- Classmates
- Medical clinic patients

## Case-control Study




15

Oregon  
Health  
Authority

15

## Case-control Study Example



16

Oregon  
Health  
Authority

16



## Advantages & Disadvantages

	Cohort	Case-control
Advantages	Can evaluate rare exposures Can calculate risk	Quick Smaller size Cost effective Can evaluate rare disease
Disadvantages	Inefficient if incubation is long Potential expense Loss to follow up	Recall bias Selection bias

## Measures of Association

- Quantifies magnitude of association between exposure and disease
  - Cohort study: relative risk (RR)
  - Case-control study: odds ratio (OR)

## 2 X 2 table

	Diseased	Not diseased	Row total
Exposed	a	b	a+b
Not exposed	c	d	c+d
Column total	a+c	b+d	a+b+c+d= Grand total

19

Oregon  
Health  
Authority

19

## Cohort Study: Relative Risk (RR)

- Risk of disease in exposed  $\div$  risk of disease in unexposed
- $RR = a/(a+b) \div c/(c+d)$

		Disease	
		Yes	No
Exposure	Yes	a	b
	No	c	d

How to interpret If  $RR = 1$ ?

20

Oregon  
Health  
Authority

20

## Calculate RR for Cohort Study

- *Salmonella* Enteritidis outbreak among group of coworkers that had catered lunch with taco bar
- 40 workers ate fresh salsa; 30 developed *Salmonella* Enteritidis and 10 did not
- Another 40 workers did not eat fresh salsa; 2 developed *Salmonella* Enteritidis and 38 did not

## Relative Risk 2 X 2 table

	<i>Salmonella</i> Enteritidis	No <i>Salmonella</i> Enteritidis	Total
Salsa	a	b	a+b
No salsa	c	d	c+d

Fill in table and calculate RR  
Put your answer in chat

## 2X2 table results

	<i>Salmonella</i> Enteritidis	No <i>Salmonella</i> Enteritidis	Total
Salsa	30	10	40
No salsa	2	38	40

$$RR = a/(a+b) \div c/(c+d)$$

$$30/(30+10) \div 2/(2+38)$$

$$RR = 15$$

## Relative Risk conclusion:

Those who ate salsa had higher risk (15X) of developing *Salmonella* Enteritidis than those who didn't eat salsa



## Case-Control Study: Odds Ratio (OR)

- Compare odds of exposure for cases and controls

- **Odds Ratio (OR) =**

➤  $a/c \div b/d$

➤  $ad/bc$

		Disease	
		Yes	No
Exposure	Yes	a	b
	No	c	d

Cross product

Oregon  
Health  
Authority

25

25

## Case-Control Study: Calculate OR

- Cases: 40 restaurant patrons with *Salmonella* Enteritidis; 30 ate eggs and 10 did not
- Controls: 40 restaurant patrons without *Salmonella* Enteritidis; 2 ate eggs and 38 did not

Why might a case-control study be good option for restaurant outbreak?

Oregon  
Health  
Authority

26

26

## Odds Ratio 2 X 2 Table

	<i>Salmonella</i> Enteritidis	No <i>Salmonella</i> Enteritidis
Eggs	a	b
No eggs	c	d
Total	a+c	b+d

Fill in table and calculate OR  
Put your answer in chat

Oregon  
**Health**  
Authority

27

27

## Odds ratio 2X2 table results

- Odds Ratio (OR) =  $(a*d)/(b*c)$
- OR=
- Conclusion:  
Cases were \_\_\_\_ times  
more likely than controls to  
have eaten eggs



Complete with your table group.

Oregon  
**Health**  
Authority

29

29

## Case-case study example: Shotgun

- Compare exposures between 2 groups of cases
- Cluster of STEC 4 cases: compare exposure frequencies with background exposure frequencies in Shotgun database
- Shotgun database provides background exposure frequencies from previously interviewed cases (Salmonella and STEC)

Exposure Code	YES	%	Backgrnd	Binomial
331 CarrotsLoose	2 / 4	50	18%	0.1547
336 PepperBell	2 / 4	50	26%	0.2813
338 PepperRed	2 / 4	50	16%	0.1180
339 PepperYellow	2 / 4	50	9%	0.0456

31

health  
Authority

31

## Is this a true association?

- P-value <0.05)  
<https://tidsskriftet.no/en/2015/09/why-p-value-significant>

- Case control study example  
OR=57 (95% CI, 12–280) P<0.0001  
<https://www.medcalc.org/calc/index.php>



32

Oregon  
Health  
Authority

32

## Study Design and Analytic Epidemiology

---

Questions?

**Stand up and stretch  
break**



Oregon  
**Health**  
Authority

33

33

## Study Design and Analytic Epidemiology

---

### Exercise

- Handout: CD 303 Exercises (pages 6-8)
- Complete each question before proceeding to next question
- Review questions together at end

Oregon  
**Health**  
Authority

34

34



## Study Design and Analytic Epidemiology

### Review of Opening Scenario

On **August 10**, the Oregon County Health Department received a report of a child with *E. coli* O157 (O157) diarrhea. This was their first O157 report in over a year. Within 24 hours Oregon County had received 3 more reports of O157 infection in children.

All 4 ill children that were reported to Oregon County Health Department over 24-hour period had attended the Oregon County Fair and had bloody diarrhea. The fair had been held **August 1 – 6**. Over ~120,000 people had attended this fair. However, the fair ended 3 days earlier. The fairground facilities are used for activities throughout the year.



35

35

## Study Design and Analytic Epidemiology

### Additional Information:

The Oregon County Fair is a major annual event in Oregon each August. It is a large agricultural fair with hundreds of animal exhibits, food vendors, and amusement rides. More than 170,000 visits were recorded this year.

The fairgrounds are supplied by shallow wells approximately 20-feet deep. Since the fairgrounds are only in operation 42 days of year, the fairground water supply is not considered a community water system but is considered an alternative water source and is regulated under the food code. This type of well must be tested within 60 days of an event. Testing for coliforms was performed on June 20<sup>th</sup> and was negative.



36

36

## Study Design and Analytic Epidemiology

---

Question 1: What hypotheses do you want to evaluate?

37

Oregon  
Health  
Authority

37

## Study Design and Analytic Epidemiology

---

### Additional information

An environmental investigation of the fairground revealed that much of the fair was supplied by chlorinated water. However, in one area of the fairground, a shallow well (well #6) provided unchlorinated water to several vendors who used the water to make beverages and ice. This area was close to the large animal barn.

38

Oregon  
Health  
Authority

38

## Study Design and Analytic Epidemiology

---

Question 2: Do we need a study, if so what type of study?

Why might case control study be used in this investigation?

39

Oregon  
Health  
Authority

39

## Study Design and Analytic Epidemiology

---

### Additional information

Cases eligible for inclusion in case control study were people who had attended the Fair and who had stool culture positive for *E. coli* O157 and were the first case in a household (primary cases).

Why limit to first case in household?

40

Oregon  
Health  
Authority

40

## Study Design and Analytic Epidemiology

---

Question 3: Whom might you get for controls? How do you find them?

41



41

## Study Design and Analytic Epidemiology

---

### Additional information

For the Oregon County Fair, controls were obtained from credit card receipts of attendees and from persons entered in photography exhibit and children's art exhibit. Controls were selected who had a similar age distribution as the cases.

42



42

## Study Design and Analytic Epidemiology

### Results

Investigators identified 46 primary cases and 84 controls. All primary cases had attended the fair during the last 4 days. Analysis included only controls who attended the fair at least once during the final 4 days and cases and controls with available exposure data (32 cases and 57 controls).

Complete 2X2 tables for each exposure.



43

43

## Study Design and Analytic Epidemiology

Question 4: What is an appropriate measure of association for a case-control study? (put in chat)

- Odds ratio
- $\frac{a/c}{b/d} = \frac{ad}{bc}$
- Measure of association for water from well #6? (put in chat)
  - OR=?
- Measure of association for Vendor A chicken? (put in chat)
  - OR=?



46

46