

Oregon Tuberculosis Control Program Indicators

1. Reporting and Treatment for TB cases

Outcome	Year																
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Completion of therapy within one year [†]	81%	88%	87%	84%	90%	94%	93%	91%	97%	97%	99%	89%	96%	87%	95%	89%	**
HIV status reported	81%	76%	77%	77%	82%	93%	78%	94%	93%	97%	96%	95%	98%	88%	99%	99%	97%
Totally directly observed therapy	55%	64%	68%	70%	61%	65%	67%	65%	84%	83%	87%	89%	80%	89%	89%	89%	**

2. Follow-up and Treatment for Contacts to Sputum-smear Positive Cases

Outcome	Year																
	2000	2001	2002	2003	2004	2005	2006*	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Cases with no contacts	2%	11%	7%	6%	7%	6%	7%	7%	0%	6%	4%	0%	0%	4%	3%	0%	0%
Contacts evaluated for TB	92%	84%	81%	88%	75%	71%	87%	80%	72%	76%	82%	74%	69%	91%	82%	74%	77%
Active TB identified	2%	2%	1%	1%	0%	1%	2%	2%	<1%	<1%	1%	0%	0%	1%	1%	0%	1%
NEW Latent TB identified [‡]	43%	39%	37%	32%	23%	19%	22%	24%	22%	18%	17%	12%	13%	17%	18%	8%	11%
LTBI treatment started [‡]	71%	93%	92%	87%	80%	76%	86%	76%	77%	84%	88%	90%	72%	84%	85%	91%	85%
LTBI treatment completed [‡]	Not Avail	85%	73%	79%	58%	69%	84%	75%	61%	71%	80%	92%	67%	70%	83%	93%	53%

Updated 11/2017

[†]Excludes patients with resistance to rifampin, patients who died during treatment, patients with meningeal TB, children under age of 15 (at report) with disseminated TB (defined as miliary and/or positive blood culture).

[‡] Excludes contacts with prior positive TST/QFT test and/or prior history of treatment for TB infection or disease.

*Oregon data from 2006 on include contact investigations to sputum smear positive AND/OR cavitary and laryngeal cases.

**Oregon data from 2016 not finalized; will be updated when complete