

**Other formats and languages**

You can get this document in other languages, large print, braille or a format you prefer free of charge. Contact the Lead Poisoning Prevention Program at [leadprogram@odhsoha.oregon.gov](mailto:leadprogram@odhsoha.oregon.gov) or 971-673-0440 (voice/text). We accept all relay calls.

## Lead Screening Protocols for Children

The goal of lead screening is to identify children who have been exposed to lead, provide appropriate interventions and reduce the risk of further exposure. If a blood lead level (BLL)  $\geq 3.5 \mu\text{g/dL}$  is detected, the nature of care and the frequency of follow-up testing vary with the patient's age and the BLL. Children with higher BLLs and their families should be educated about what lead exposure is and what they can do about it. The single most important factor in managing childhood lead exposure is identifying and reducing the child's exposure to lead. For more guidance on caring for children with higher blood lead levels see the website [Management of Childhood Lead Exposure](#) from PEHSU.

### Anticipatory Guidance

Recent research has shown that there is no safe level of lead exposure and levels below the [blood lead reference value](#) ( $3.5 \mu\text{g/dL}$ ) can cause permanent problems with learning, memory and behavior. Therefore, anticipatory guidance regarding lead hazard identification and risk reduction measures should be a routine part of an ongoing educational approach for pregnant people, children and their families. Health Care Providers should provide source identification and risk reduction educational materials.

Lead exposure during pregnancy is especially problematic as lead can cross the placenta and interfere with normal development of the fetal brain. Pregnant people can be exposed to lead through a variety of sources. People who are pregnant or likely to become pregnant should try to maintain lead levels below  $3.5 \mu\text{g/dL}$  and as low as possible. Anticipatory guidance should focus on decreasing the risk of exposure to lead by advising against activities such as remodeling or repainting the baby's room or restoring old furniture.

## Testing Methods

Blood lead testing is the only acceptable laboratory test for screening and confirming lead poisoning. Venipuncture is preferred for specimen collection, but capillary testing is acceptable if care is taken to properly clean and prepare the finger. Capillary samples are easier to contaminate because of the possibility of lead containing dust and dirt on the hand or under the fingernails. All capillary BLLs of 3.5 µg/dL or higher must be followed with a confirmatory venous test. The higher the capillary screening BLL, the more urgent the need for a venous confirmatory test. The schedule for confirmatory testing of a child with an EBLL is in table 3. Several tests have been found to be insensitive and/or imprecise as screening tests for lead and are not recommended. These include erythrocyte protoporphyrin (EP), zinc protoporphyrin (ZPP) basophilic stippling, urine testing, and assays of hair or fingernail lead levels.

## Oregon Lead Risk Assessment Questionnaire

All children should be assessed for risk of lead poisoning by administration of the Oregon Lead Risk Assessment Questionnaire (see below). This questionnaire should be administered at 12 and 24 months of age, or between 3 and 5 years of age if not previously screened. If the answer to any of the following questions is “Yes” or “Don’t know” a blood lead test should be performed. Follow-up questions may be needed to clarify responses.

Per [OAR 410-130-0246](#), all children on Medicaid/Oregon Health Plan must be blood tested at 12 and 24 months, or between 24 and 72 months if there is no record of a previous blood test.

1. Has your child lived in or regularly visited a home, child care or other building built before 1950?
2. Has your child lived in or regularly visited a home, child care or other building built before 1978 with recent or ongoing painting, repair and/or remodeling?
3. Is your child enrolled in or attending a Head Start program or receiving WIC benefits *with no record of a previous blood lead test??*
4. Does your child have a brother, sister, other relative, housemate or playmate with lead poisoning?
5. Does your child spend time with anyone that has a job or hobby where they may work with lead?

*Examples: painting, remodeling, auto radiators, batteries, auto repair, soldering, making sinkers, bullets, stained glass, pottery, going to shooting ranges, hunting or fishing.*

6. Do you have pottery or ceramics (*barro*) made in Mexico/Latin/South America that are used for cooking, storing or serving food or drink?
7. Has your child ever taken any traditional home remedies or used imported cosmetics?  
*Examples: Azarcon/Alarcon/Greda, Kohl/Surma/Kajal, Sindoor/Tikka, Ayurvedic*
8. Has your child been adopted from, lived in or visited another country in the past 6-12 months?
9. Do you have concerns about your child's development or behavior?

## Diagnostic Blood Lead Testing

Blood lead testing should also be considered as part of a diagnostic work-up of any child regardless of age with the following symptoms:

- **Behavioral problems:** aggression, hyperactivity, attention deficit, school problems, learning disabilities, excessive mouthing or pica behavior and other behavior disorders.
- **Developmental problems:** growth, speech and language delays and/or hearing loss.
- **Symptoms or signs consistent with lead poisoning:** irritability, headaches, vomiting, seizures or other neurological symptoms, anemia, loss of appetite, abdominal pain and cramping or constipation.
- **Ingestion of foreign objects:** paint chips, jewelry, fishing weight, ammo
- **Proximity to airport:** exposure to lead in air and soil from aviation gas used in piston engine aircraft.

## Confirmatory Testing Schedule

Any capillary screening BLL  $\geq 3.5$   $\mu\text{g/dL}$  should be confirmed with a venous sample. The higher the BLL on the capillary test, the more urgent the need for confirmatory testing. The table below outlines the confirmatory venous schedule based on the initial screening result.

BLL ( $\mu\text{g/dL}$ )	Confirmation Testing (venous)	Follow-Up Testing (venous)
3.5-9	As soon as possible, or within 7-14 days	3 months
10-19	As soon as possible, or within 7 days	3 months

<b>BLL (µg/dL)</b>	<b>Confirmation Testing (venous)</b>	<b>Follow-Up Testing (venous)</b>
<b>20-44</b>	As soon as possible, or within 7 days	1 month
<b>45-59</b>	As soon as possible, or within 2 days	Chelation* with subsequent follow-up.
<b>60-69</b>	As soon as possible, or within 1 day	Chelation* with subsequent follow-up
<b>≥ 70</b>	Immediately as an emergency lab test	Chelation* with subsequent follow-up

**Exception to confirmatory testing schedule:** If recent known exposure (e.g. foreign body ingestion, recent remodeling, etc.) confirm as soon as possible for all blood lead levels.

\*While chelation therapy is considered a mainstay in the medical management of children with BLLs  $\geq 45$  µg/dL, it should be used with caution. Unless the clinician is intimately familiar with treatment protocols, he/she should consult with an expert such as one of the following prior to using chelation agents:

- Medical toxicologist
- Clinician experienced in treating children with elevated BLLs
- [Northwest Pediatric Environmental Specialty Health Unit \(PESHU\)](#)
- Center with expertise in the management of lead chemotherapy. Clinicians may contact the [Oregon Poison Center](#) for the names of accessible experts.

## Public Health Division

Lead Poisoning Prevention Program  
 800 NE Oregon St. Suite 640  
 Portland, OR 97232  
 971-673-0440  
[leadprogram@odhsoha.oregon.gov](mailto:leadprogram@odhsoha.oregon.gov)  
[healthoregon.org/lead](http://healthoregon.org/lead)

