# Maternal Depression Screening & Referrals Overview for MIECHV EHS/HFA/NFP Programs



September 2020



1

## Objectives



#### **Short-term**

Increase understanding of when and how to complete and document screening and referrals for Maternal Depression



#### Medium-term

Demonstrate through data the incredible work home visitors engage in to provide Maternal Depression screening and referrals to clients they serve



#### Long-term

Continue to maintain or increase MIECHV funding for home visiting services in Oregon




### MIECHV data collection

- Required for all MIECHV grantees
- Demonstrates the work of MIECHV at a national level
- Supports quality improvement efforts
- Demonstrate measurable improvement over time



3





	Measure Definition	FY 2019
		78.5%
Performance Measure #3: Maternal Depression Screening	Percent of primary caregivers who are screened for depression using a validated tool within three months of enrollment (for those not enrolled prenatally) or within three months of delivery (for those enrolled prenatally).	332/423
	Bright spot: this measure increased from 2018 (56.7%)  Herealth	

## When do maternal depression screenings occur for MIECHV?

Within 90 days of enrollment (for those not enrolled prenatally)

OR

Within 90 days of delivery (for those enrolled prenatally)

→ See client schedule for window period; if screened before or after this period, doesn't count for measure



H

7

#### What is required for maternal depression screening?


- All clients should receive a PHQ-9 screening regardless of whether they are currently or have previously received mental health services
- A PHQ-9 screen must be offered and completed to meet the measure.


Maternal & Child Health Section Public Health Division



#### Maternal Depression Screening Data Collection form and THEO entry for EHS HFA programs: M5 MIECHV Child's Enrollment Tool Part A: Depression Screening To meet the measure, all clients must receive a PHQ-9 screening within 90 days of enrollmer infant birth regardless of whether they are currently receiving or have previously received depres-services. M5 - MIECHV Child's Enrollment Tool - Index Parent\_20201001 1. PHQ-9 completed? $\square$ Yes, completed $\rightarrow$ Date tool completed: $\underline{\hspace{1cm}/\hspace{1cm}}$ / 20 $\underline{\hspace{1cm}}$ $\rightarrow$ Go to Question 1a. 1a. If Yes, result of PHQ-9: Score of 10 or higher → Go to Part B: Depression Referral. Score of 9 or lower No, not completed □ No, not completed → Complete Screening at future visit before the 90-day window close complete another M5 form and update in THEO. Go to Question 1a. To meet the measure, client must receive a Referral AND at least one Service if the client so 10 or higher on the PHQ-9 during a MIECHV screening and is not currently receiving depression 1a. If Yes, result of PHQ-9: 2. If a Score of 10 or higher, did you provide a referral? \*See examples below Score of 9 or lower Yes → Complete M2B-MIECHV Referral Tracking & Follow-up Form. ☐ No, the client is currently receiving services (Clients who are currently receiving depred on not need to be referred). 2. If a Score of 10 or higher, did you provide a referral? ☐ No, other reason → Clients who screen positive should be offered and start/receive at length referral service. Continue to follow up with client at future visits to track referral and follow-up M2B-MIECHV Referral Tracking & Follow-up Form. No. the client is currently receiving services No, other reason \*Recommended referral services include: Internal referral to self or other staff trained in Mothers and Babies intervention; Licensed or licensed-eligible mental health provider (LMHC, LCSW, or LMFT); Medication therapy from a primary care provider, psychiatrist, or women's health provider.

9





Tips for completing maternal depression screening

If you are unable to complete the screening during the visit, try and complete during the next one

11

11

## PHQ-9 and Maternal Depression Screening training and resources:

https://www.oregon.gov/oha/PH/HEALTHYPEOPLEF AMILIES/BABIES/HOMEVISITING/MIECHV/Pages/mie chv-orientation.aspx (scroll down webpage)

#### Remote maternal depression screening resources:

<u>Screenings in Virtual Visits</u> created by <u>the Rapid</u> <u>Response-Virtual Home Visiting Collaborative</u>

IPV HV CollN memo: <u>Coronavirus/COVID-19 and Implications for Maternal Depression and Intimate</u> Partner Violence Screening and Referral



Maternal
Depression
Screening
Resources

	Measure Definition	FY 2019
Performance Measure #17: Maternal Depression Referrals	Percent of primary caregivers referred to services for a positive screen for depression who receive one or more service contacts.	<b>22.7%</b> (15/66)
	Bright spot: this measure increased from 2018!	

When are maternal depression referrals needed?

 If the client scores a 10 or higher on the PHQ-9 during the MIECHV screening <u>and</u> is not currently receiving depression services

If the client receives a referral and then receives referral service:

 Update the M2B-MIECHV Referral Tracking and Follow-up form in THEO (for EHS/HFA) or update and resubmit the form to OHA (for NFP)



14

## Considerations for meeting the maternal depression measure





Caregivers who screen positive for maternal depression will remain in the measure denominator until they receive a service ("referral contact").



There is no timeframe for when the client must receive a referral contact  $\rightarrow$  continue to follow up with client to support them in accessing service


15


## What Counts as a Depression Referral??

Caregiver must receive a referral and at least one service

#### · Recommended referral services include:

- Internal referral to self or other staff member trained in Mothers and Babies intervention
- Internal referral to self or other staff member trained in Mental Health Intervention (for NFP programs)
- Licensed or licensed-eligible mental health provider (LMHC, LCSW, or LMFT) using cognitive behavioral therapy or another evidence-based treatment
- Medication therapy from a primary care provider, psychiatrist, or women's health provider





### Thank You!

For more information about Oregon MIECHV Program's data collection, please contact:

**Tina Kent**, MIECHV Data Manager: <a href="mailto:TINA.M.KENT@dhsoha.state.or.us">TINA.M.KENT@dhsoha.state.or.us</a>
<a href="mailto:Drewallyn.b.riley@dhsoha.state.or.us">Drewallyn Riley</a>, CQI Coordinator: <a href="mailto:Drewallyn.b.riley@dhsoha.state.or.us">Drewallyn.b.riley@dhsoha.state.or.us</a>
<a href="mailto:For THEO Application Support: theo.support@state.or.us">Drewallyn.b.riley@dhsoha.state.or.us</a>
<a href="mailto:For THEO Application Support: theo.support@state.or.us">THEO Application Support: theo.support@state.or.us</a>

Health

Public Health Division Maternal & Child Health