Safe Sleep for Babies



Always place your baby on his or her back for naps and at bedtime.



Facts about SIDS

(Sudden Infant Death Syndrome)

- In Oregon, about 40 babies die in their sleep every year.
- The most important thing you can do for your baby's health is to quit smoking. For help, call the Oregon Tobacco Quit Line at 1-800-784-8669 or online at www.quitnow.net/oregon.
- Breastfeeding reduces your baby's risk of SIDS.

Some parents choose to sleep with their baby. It can be unsafe to sleep with your baby if:

- Your baby is less than 12 weeks old.
- Your bed has pillows, quilts, or comforters.
- You share the bed with pets or other children.
- You or your partner:
 - Smokes
 - Has taken drugs, alcohol, or medications that make you sleepy
 - · Are very tired or suffer from extreme exhaustion.

To make sure your baby is as safe as possible:

- Put your baby to sleep on his or her back.
- Make sure no one smokes around your baby.
- Breastfeed your baby.
- Put your baby to sleep in the room where you sleep.
- Remove soft fluffy bedding, pillows, stuffed toys and bumper pads from your baby's sleep area.
- Face up Face clear Smoke-free Baby near!
- Avoid placing your baby to sleep on soft things like couches or upholstered chairs, waterbeds, beanbag chairs, quilts or comforters.
- Talk with grandparents and other caregivers about this safe sleep information.

Please talk to your health care provider about any concerns you may have.

Contact for more information 971-673-0262 Visit Babies First! Safe Sleep for Babies at www.healthoregon.org/safesleep.

This document can be provided upon request in alternative formats for individuals with disabilities. Other formats may include (but are not limited to) large print, Braille, audio recordings, Web-based communications and other electronic formats. Call 971-673-0252 (voice) or 971-673-0372 (TTY) to arrange for the alternative format that will work best for you.