

Title:RSV season has ended in Oregon and clinicians may discontinue offering RSV prevention products

On May 2, Oregon met criteria for respiratory syncytial virus (RSV) season offset. While RSV continues to circulate at very low levels, clinicians may discontinue offering infant immunizations (nirsevimab/clesrovimab) and the maternal RSV vaccine (Abrysvo).

Because RSV seasonality is unpredictable in Oregon, Oregon Health Authority (OHA) uses local epidemiology to guide RSV immunization recommendations. OHA will continue to send HANs alerting providers to the onset and offset of each RSV season.

Oregon's RSV Surveillance Report is available here:

[https://www.oregon.gov/oha/PH/DISEASES/CONDITIONS/COMMUNICABLEDISEASE/DISEASESURVEILLANCEDATA/Documents/RSV/RSV\\_Oregon.pdf](https://www.oregon.gov/oha/PH/DISEASES/CONDITIONS/COMMUNICABLEDISEASE/DISEASESURVEILLANCEDATA/Documents/RSV/RSV_Oregon.pdf)

Additional information regarding RSV immunizations for infants and young children may be found here: <https://www.cdc.gov/rsv/hcp/vaccine-clinical-guidance/infants-young-children.html>

For questions, please contact Melissa Sutton at [Melissa.Sutton@oha.oregon.gov](mailto:Melissa.Sutton@oha.oregon.gov).