

## Wheeler County: Adult Immunization Rates

	2016-2017	2017-2018	2018-2019	2019-2020
<b>Influenza Vaccination Rates<sup>a,b</sup></b>				
<b>Female</b>				
18 to 49 years	37.5%	35.1%	34.7%	39.2%
50 to 64 years	49.8%	46.3%	54.8%	58.3%
≥65 years	51.4%	57.6%	59.1%	62.6%
<b>Male</b>				
18 to 49 years	13.0%	16.1%	19.0%	20.6%
50 to 64 years	35.1%	34.5%	36.3%	36.0%
≥65 years	46.3%	53.1%	51.3%	58.3%
<b>All adults</b>				
18 to 49 years	25.7%	25.8%	26.2%	29.9%
50 to 64 years	43.9%	41.6%	47.3%	48.8%
≥65 years	49.0%	55.4%	55.4%	60.5%
		<b>2017</b>	<b>2018</b>	<b>2019</b>
<b>Tetanus (Td or Tdap) Vaccination Rates<sup>b,c</sup></b>				
<b>All adults</b>				
≥18 years		5.7%	6.2%	5.5%
18 to 49 years		6.6%	6.4%	7.6%
50 to 64 years		3.9%	6.4%	6.0%
≥65 years		6.2%	6.0%	4.1%
<b>Zoster (Shingles) Vaccination Rates<sup>b,d</sup></b>				
<b>All adults</b>				
50 to 59 years		0.5%	6.7%	8.0%
≥60 years		4.9%	5.2%	12.9%
60 to 64 years		7.5%	10.0%	18.2%
≥65 years		4.1%	4.0%	10.7%

<sup>a</sup> Immunization rates reflect stated flu season

<sup>b</sup> Estimates are based on a combination of ALERT Immunization Information System (IIS, Oregon Immunization Program) and Portland State University's Population Research Center data for Oregon populations

<sup>c</sup> Target immunization rate for the overall adult population is 10% per year; however, target rates are higher for women of child-bearing age

<sup>d</sup> Rate reflects vaccination with either Shingrix or Zostavax