

Oregon Region 2 Pediatric Readiness Quality Improvement Report

Region: Region 2

Number of Emergency Departments (EDs) Responding in Region: 10

Report Date: 2/8/2017 8:12:53 PM

Region 2
Average Score
Out of 100

72

Oregon
Average Score
Out of 100

68

About Pediatric Readiness:

The National Pediatric Readiness Project is a national multi-phase quality improvement initiative to ensure all U.S. emergency departments have the essential guidelines and resources in place to provide effective emergency care to children. The total number of points possible is 100. A score of 100 represents the essential components needed to establish a foundation for pediatric readiness, but is in no way inclusive of all the components recommended for pediatric readiness.

Pediatric Patient Volume and Scores for Region 2:

Pediatric Volume	Number EDs in Region	Minimum Score	Maximum Score	Average Score
High Volume (>10,000 patients)	1	*	*	*
Medium High Volume (5,000 to 9,999 patients)	1	*	*	*
Medium Volume (1,800 to 4,999 patients)	5	44	82	61
Low Volume (<1,800 patients)	3	*	*	*

* Indicates the number of EDs is less than 5 so scores cannot be displayed. This asterisk will be shown on the following pages when applicable.

Region 2

Pediatric Readiness Overall Section Scoring:

Section	QI Status	% of Region EDs Have	Avg. Region Score	Avg. Oregon Score	Section Point Value
Guidelines for Administration and Coordination (19 pts)		45.0%	8.6	8.5	19
Physicians, Nurses, and Other Health Care Providers Who Staff the ED (10 pts)		55.0%	5.5	5.0	10
Guidelines for QI/PI in the ED (7 pts)		54.3%	3.8	2.5	7
Guidelines for Improving Pediatric Patient Safety in the ED (14 pts)		83.5%	11.7	11.8	14
Guidelines for Policies, Procedures, and Protocols for the ED (17 pts)		70.4%	12.0	10.5	17
Guidelines for Equipment, Supplies, and Medications for the Care of Pediatric Patients in the ED (33 pts)		90.9%	30.0	29.6	33

QI Status:

-  All EDs Have
-  80 to 90% Have
-  60 to 79% Have
-  30 to 59% Have
-  Less Than 29% Have

Region 2

Guidelines for Administration and Coordination of the ED for the Care of Children:

Question	QI Status	Number EDs Have	% of Region EDs Have	% of State EDs Have	Question Point Value
Physician Coordinator	*			45.5%	9.5
Nurse Coordinator	◆	5	50.0%	43.6%	9.5

Physicians, Nurses, and Other Health Care Providers Who Staff the ED:

Physician Competency Evaluations	*			32.7%	5.0
Nurse Competency Evaluations	●	8	80.0%	67.3%	5.0

Guidelines QI/PI in the ED:

Does your ED have a pediatric patient care-review process?	▲	6	60.0%	38.2%	5.0
Identification of quality indicators for children	*			27.3%	0.5
Collection and analysis of pediatric emergency care data	◆	5	50.0%	36.4%	0.5
Development of a plan for improvement in pediatric emergency	*			30.9%	0.5
Re-evaluation of performance using outcomes-based measures	*			23.6%	0.5

QI Status:

 All EDs Have
  80 to 90% Have
  60 to 79% Have
  30 to 59% Have
  Less Than 29% Have

Pediatric Patient Safety in the ED:

Question	QI Status	Number EDs Have	% of Region EDs Have	% of State EDs Have	Question Point Value
Weigh and Record in Kg		6	60.0%	70.9%	3.5
Temperature, heart rate, and respiratory rate recorded		10	100.0%	96.4%	1.4
Blood pressure monitoring available based on severity of illness		10	100.0%	100.0%	1.4
Pulse oximetry monitoring available based on severity of illness		10	100.0%	100.0%	1.4
Written procedure in place for notification of physicians when abnormal vital signs		6	60.0%	74.5%	1.4
Process in place for the use of pre-calculated drug dosing		9	90.0%	80.0%	3.5
Process in place that allows for 24/7 access to interpreter services in the ED		10	100.0%	98.2%	1.4

QI Status:

-  All EDs Have
-  80 to 90% Have
-  60 to 79% Have
-  30 to 59% Have
-  Less Than 29% Have

Region 2

Policies and Procedures:

Question	QI Status	Number EDs Have	% of Region EDs Have	% of State EDs Have	Question Point Value
Triage policy that specifically addresses ill and injured children		7	70.0%	65.5%	2.12
Policy for pediatric patient assessment and reassessment		8	80.0%	74.5%	1.70
Policy for immunization assessment and management of the under-immunized child		6	60.0%	52.7%	1.70
Policy for child maltreatment		9	90.0%	85.5%	1.70
Policy for death of the child in the ED		7	70.0%	52.7%	1.70
Policy for reduced-dose radiation for CT and x-ray imaging based on pediatric age or weight		8	80.0%	63.6%	1.70
Policy for promoting family-centered care		7	70.0%	65.5%	2.12
Hospital disaster plan addresses issues specific to the care of children		5	50.0%	41.8%	2.12
Inter-facility transfer guidelines		7	70.0%	58.2%	2.12

QI Status:

 All EDs Have
  80 to 90% Have
  60 to 79% Have
  30 to 59% Have
  Less Than 29% Have

Region 2

Equipment and Supplies:

Availability, Location, Training

Question	QI Status	Number EDs Have	% of Region EDs Have	% of State EDs Have	Question Point Value
Is the ED staff trained on the location of all pediatric equipment and medications?		10	100.0%	98.2%	1.0
Is there a daily method used to verify the proper location and function of pediatric equipment and supplies?		9	90.0%	87.3%	1.0
Is a medication chart, length-based tape, medical software, or other system readily available to ensure proper sizing of resuscitation equipment and proper dosing of medications?		10	100.0%	100.0%	1.0

QI Status:

-  All EDs Have
-  80 to 90% Have
-  60 to 79% Have
-  30 to 59% Have
-  Less Than 29% Have

Region 2

Equipment and Supplies: Monitoring Equipment Items

Question	QI Status	Number EDs Have	% of Region EDs Have	% of State EDs Have	Question Point Value
Neonatal blood pressure cuff	✓	10	100.0%	96.4%	0.55
Infant blood pressure cuff	✓	10	100.0%	100.0%	0.55
Child blood pressure cuff	✓	10	100.0%	100.0%	0.55
Defibrillator with pediatric and adult capabilities including pads/paddle	✓	10	100.0%	100.0%	0.55
Pulse oximeter with pediatric and adult probes	✓	10	100.0%	100.0%	0.55
Continuous end-tidal CO2 monitoring device	✓	10	100.0%	100.0%	0.55

QI Status:

- ✓ All EDs Have
- 80 to 90% Have
- ▲ 60 to 79% Have
- ◆ 30 to 59% Have
- ✗ Less Than 29% Have

Region 2

Equipment and Supplies: Fluid Resuscitation Equipment Items

Question	QI Status	Number EDs Have	% of Region EDs Have	% of State EDs Have	Question Point Value
22 gauge catheter-over-the-needle		10	100.0%	100.0%	0.55
24 gauge catheter-over-the-needle		10	100.0%	100.0%	0.55
Pediatric intra-osseus needles		10	100.0%	100.0%	0.55
IV administration sets with calibrated chambers and extension tubing and/or infusion devices with ability to regulate rate and volume of infusate		10	100.0%	96.4%	0.55
Umbilical vein catheters (3.5F or 5.0F)		8	80.0%	61.8%	0.55
Central venous catheters (any two sizes in range, 4F-7F)		5	50.0%	52.7%	0.55

QI Status:

 All EDs Have
  80 to 90% Have
  60 to 79% Have
  30 to 59% Have
  Less Than 29% Have

Region 2

Equipment and Supplies:

Respiratory/Airway Management Equipment Items

Question	QI Status	Number EDs Have	% of Region EDs Have	% of State EDs Have	Question Point Value
Endotracheal tubes: cuffed or uncuffed 2.5 mm	✓	10	100.0%	92.7%	0.557
Endotracheal tubes: cuffed or uncuffed 3.0 mm	✓	10	100.0%	98.2%	0.557
Endotracheal tubes: cuffed or uncuffed 3.5 mm	✓	10	100.0%	100.0%	0.557
Endotracheal tubes: cuffed or uncuffed 4.0 mm	✓	10	100.0%	100.0%	0.557
Endotracheal tubes: cuffed or uncuffed 4.5 mm	✓	10	100.0%	100.0%	0.557
Endotracheal tubes: cuffed or uncuffed 5.0 mm	✓	10	100.0%	100.0%	0.557
Endotracheal tubes: cuffed or uncuffed 5.5 mm	✓	10	100.0%	96.4%	0.557
Endotracheal tubes: cuffed or uncuffed 6.0 mm	✓	10	100.0%	100.0%	0.557
Laryngoscope blades: straight, size 00	●	8	80.0%	74.5%	0.557
Laryngoscope blades: straight, size 0	✓	10	100.0%	92.7%	0.557
Laryngoscope blades: straight, size 1	✓	10	100.0%	98.2%	0.557
Laryngoscope blades: straight, size 2	✓	10	100.0%	96.4%	0.557
Laryngoscope blades: curved, size 2	✓	10	100.0%	98.2%	0.557
Pediatric-sized Magill forcep	●	9	90.0%	85.5%	0.557

QI Status:

 All EDs Have
  80 to 90% Have
  60 to 79% Have
  30 to 59% Have
  Less Than 29% Have

Region 2

Equipment and Supplies:

Respiratory/Airway Management Equipment Items (Cont. 2)

Question	QI Status	Number EDs Have	% of Region EDs Have	% of State EDs Have	Question Point Value
Nasopharyngeal airways: infant-size		8	80.0%	78.2%	0.557
Nasopharyngeal airways: child-size		9	90.0%	83.6%	0.557
Oropharyngeal airways: size 0 (50mm)		10	100.0%	96.4%	0.557
Oropharyngeal airways: size 1 (60mm)		10	100.0%	94.5%	0.557
Oropharyngeal airways: size 2 (70mm)		10	100.0%	96.4%	0.557
Oropharyngeal airways: size 3 (80mm)		10	100.0%	94.5%	0.557
Stylets for pediatric/infant-sized endotracheal tube		10	100.0%	96.4%	0.557
Tracheostomy tubes: size 3.0 mm		5	50.0%	54.5%	0.557
Tracheostomy tubes: size 3.5 mm		5	50.0%	56.4%	0.557
Tracheostomy tubes: size 4.0 mm		5	50.0%	58.2%	0.557
Bag-mask device, self inflating: infant, 450 ml		10	100.0%	96.4%	0.557
Masks to fit bag-mask device adaptor: neonate		9	90.0%	90.9%	0.557

QI Status:

All EDs Have
 80 to 90% Have
 60 to 79% Have
 30 to 59% Have
 Less Than 29% Have

Region 2

Equipment and Supplies:

Respiratory/Airway Management Equipment Items (Cont. 3)

Question	QI Status	Number EDs Have	% of Region EDs Have	% of State EDs Have	Question Point Value
Masks to fit bag-mask device adaptor: infant	●	9	90.0%	92.7%	0.557
Masks to fit bag-mask device adaptor: child	●	9	90.0%	92.7%	0.557
Clear oxygen masks: standard infant	✓	10	100.0%	90.9%	0.557
Clear oxygen masks: standard child	✓	10	100.0%	100.0%	0.557
Non-rebreather masks: infant-sized	●	8	80.0%	85.5%	0.557
Non-rebreather masks: child-size	✓	10	100.0%	94.5%	0.557
Nasal cannulas: infant	✓	10	100.0%	89.1%	0.557
Nasal cannulas: child	✓	10	100.0%	94.5%	0.557
Laryngeal mask airways: size 1	▲	6	60.0%	70.9%	0.557
Laryngeal mask airways: size: 1.5	▲	6	60.0%	65.5%	0.557
Laryngeal mask airways: size: 2	▲	7	70.0%	74.5%	0.557
Laryngeal mask airways: size: 2.5	▲	7	70.0%	70.9%	0.557
Laryngeal mask airways: size: 3	▲	7	70.0%	81.8%	0.557

QI Status:

 All EDs Have
  80 to 90% Have
  60 to 79% Have
  30 to 59% Have
  Less Than 29% Have

Region 2

Equipment and Supplies:

Respiratory/Airway Management Equipment Items (Cont. 4)

Question	QI Status	Number EDs Have	% of Region EDs Have	% of State EDs Have	Question Point Value
Suction catheters: at least one in range 6-8F		10	100.0%	98.2%	0.557
Suction catheters: at least one in range 10-12F		10	100.0%	100.0%	0.557
Supplies/kit for pediatric patients with difficult airways (supraglottic airways of all sizes, needle cricothyrotomy supplies, surgical cricothyrotomy kit)		8	80.0%	83.6%	0.557

QI Status:

-  All EDs Have
-  80 to 90% Have
-  60 to 79% Have
-  30 to 59% Have
-  Less Than 29% Have