

# Randall Children's Hospital at Legacy Emanuel

## Pain control options for venipunctures and IV placement

| Instructions/Contraindications                        |   | Age Consideration  |
|---|---|--|
| <b>Lidocaine (ELA-MAX) 4% cream, 5g</b>               | <ul style="list-style-type: none"> <li>• Must be placed approximately 30 minutes prior to poke for peak effect</li> <li>• Tip: Rub a small amount of ELA-MAX into the skin before placing the desired amount with tegaderm</li> <li>• Clean site prior to poke</li> <li>• Contraindicated in patients &lt; 37 weeks gestation and/or those with a lidocaine allergy</li> </ul>  | ≥ 2 months   |
| <b>Lidocaine 1% for use with J-tip syringe 0.2 ml</b> | <ul style="list-style-type: none"> <li>• Explain "rocket-ship" or soda can opening sound to patient                             <ul style="list-style-type: none"> <li>◦ Child Life has an mp3 of the sound it makes and can assist with medical play</li> </ul> </li> <li>• Manipulate the skin to avoid injecting directly over the vein</li> <li>• Hold J-tip against the skin at a 90° angle before releasing.</li> <li>• Clean after using the J-tip and prior to poke. (see tip sheet)</li> <li>• Contraindicated for patients:                             <ul style="list-style-type: none"> <li>◦ Receiving chemotherapy or blood thinners</li> <li>◦ Bleeding disorder</li> <li>◦ &lt; 2 months of age</li> <li>◦ Needing port access</li> <li>◦ Lidocaine allergy</li> </ul> </li> </ul> <p style="text-align: right;">demo click here: <a href="#">J-Tip Demo</a></p> | Can be used >2mo of age<br><br>Preschoolers or younger child/infant may find the noise frightening, ensure preparation for all ages. |
| <b>Ethyl Chloride spray 100%</b>                      | <ul style="list-style-type: none"> <li>• Clean skin, spray site continuously for 3-7 seconds (&lt;10 seconds) from a distance of 3-9 inches away</li> <li>• Effects last up to 60 seconds, spray immediately prior to the procedure (see tip sheet)</li> <li>• Contraindicated on open wounds, mucus membranes, and/or for children &lt; 3 years old</li> </ul> <p style="text-align: right;">demo (~00:35-1:35) click here: <a href="#">Ethyl Chloride Demo</a></p>  | ≥ 2 years  |
| <b>Sucrose</b>  | <ul style="list-style-type: none"> <li>• Provide drops PO immediately prior to and during procedure</li> </ul>  | 0 to 12 months   |
| <b>Shot blocker</b>                                   | <ul style="list-style-type: none"> <li>• For injections i.e immunizations or flu shots (single use-may give to caregiver to use at PCP)</li> <li>• Clean site prior to use, position and hold firmly against skin during injection</li> </ul>   | All ages   |

### Ordering Steps:

- |  |                                       |  |
|--|---------------------------------------|--|
| 1. Click on "orders" then "go to order sets" | 4. Click "open order sets"            | (LIP Sig Required)"                              |
| 2. In the search box type "topical"          | 5. Click the box of the desired order | 8. Select the attending provider, click "accept" |
| 3. Right click to "add to favorites"         | 6. Click "sign"                       | 9. Scan medication prior to use                  |
|  | 7. Order mode: "Protocol Initiation"  |  |

