Table III OAR 333-077-0125

Condition of <u>NEWBORN AFTER DELIVERY</u> that Requires Transfer or Consultation

Conditions highlighted in red boxes indicate high-risk conditions that, when condition develops, will require the transfer of a client to hospital-based care.

Conditions in the yellow boxes indicate potentially risky conditions that require consultation with a provider of perinatal care as defined under OAR 333-077-0125.

The list of high-risk conditions and potentially risky conditions is not comprehensive, and other physical, behavioral health, obstetric, or fetal conditions may arise that require consultation or transfer to hospital-based care. Having multiple risk conditions requiring consultation may increase the risk sufficiently to indicate the need to transfer care.

Apgar	• Low Apgar score < 5 at 5 minutes or < 7 at 10 minutes
Birth injury	Enlarging cephalohematoma
	Significant birth injury
	Excessive bruising
Blood glucose	Hyperglycemia unresponsive to treatment
	Hypoglycemia unresponsive to treatment
Congenital anomaly	Unexpected and life-threatening
	Expected and documented, life-threatening
Gastrointestinal	Emesis of fresh blood
	Persistent projectile or bilious vomiting
Infection	Evident or suspected infection
	• Persistent inability to maintain temperature between 97 to 100 degrees Fahrenheit or 36 to 37 degrees Celsius.
Neurological	Persistent, unexplained hypotoniaSeizures
Respiratory	Central Cyanosis Unresolved Pallor
	 Heart rate less than 80 or greater than 160 (at rest) without improvement.
	Persistent cardiac murmur
	Respiration rate greater than 100 within the first two bours postportum, and greater than 20 thereafter.
	hours postpartum, and greater than 80 thereafter, lasting more than one hour without improvement.
Weight	Weight less than 2,270 grams (five pounds)

Effective date: November 3, 2025