

Unadjusted screening among adults by race and ethnicity, Oregon 2015-2017

Use this table to estimate the overall burden of disease within each race and ethnicity



Screening type		White, NL	African American, NL	American Indian or Alaska Native, NL	Asian, NL	Pacific Islander, NL	Asian or Pacific Islander, NL	Latino/a
Mammogram within two years ¹	Unadjusted (%)	76.4	69.8	61.9	85.8	--	82.8	74.7
	Number of adults	426,300	6,100	8,600	22,100	--	23,300	102,400
Pap test within three years ²	Unadjusted (%)	81.0	86.1	85.6	78.9	61.1	76.0	77.5
	Number of adults	452,000	7,500	25,700	61,200	5,300	66,500	106,200
Stool (FOBT) test within past year ³	Unadjusted (%)	13.1	20.9	15.5	15.3^	--	15.4^	10.9
	Number of adults	73,100	1,800	4,000	7,000	--	7,800	7,500
Colorectal cancer screening ⁴	Unadjusted (%)	71.5	63.8	55.6	67.2	--	65.0	60.9
	Number of adults	398,900	5,500	14,400	30,900	--	32,800	41,900
Cholesterol check within the past five years	Unadjusted (%)	81.0	72.4	75.8	67.1	61.1	66.0	71.5
	Number of adults	2,138,500	51,800	55,200	114,600	12,200	126,800	373,600

NL = non-Latino

^ This number may be statistically unreliable and should be interpreted with caution.

-- This number is suppressed because it is statistically unreliable.

1. Women age 50-74.

2. Women age 21-65.

3. Men and women age 50-74.

4. Colonoscopy within 10 years; Sigmoid within the past 5 years with fecal occult blood test in past 3; FOBT in the last year; among men and women age 50-75.

Source: Oregon Behavioral Risk Factors Surveillance System Race Oversample Dataset 2015-2017.

Suggested citation: Oregon Health Authority, Public Health Division, Health Promotion and Chronic Disease Prevention section. Screening among adults by race and ethnicity, Oregon 2015-2017. <https://www.oregon.gov/oha/PH/DISEASES/CONDITIONS/CHRONICDISEASE/DATAREPORTS/Pages/index.aspx>

Created April 19, 2019. Accessed [DOWNLOAD DATE].