



	0 – 6 months	6 months –1 year	1 – 5 years	5-9 years	9-11 years	11-18 years	18-19 years	19-27 years	27-45 years	45-50 years	50-65 years	65 years +
DTaP, DT, Tdap, Td	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal
COVID-19		Up to 23 months	Some	Some	Some	Some	Some	Some	Some	Some	Some	Some
Hep. A		Some	Some	Some	Some	Some	Some	Some	Some	Some	Some	Some
Hep. B	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal
HIB	Universal	Universal	Universal	Some	Some	Some	Some	Some	Some	Some	Some	Some
HPV					Some	Some	Some	Some	Some	Some		
Influenza		Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal
MMR		Some	Some	Some	Some	Some	Some	Some	Some	Some	Some	Some
Mening. ACYW	Some	Some	Some	Some	Some	Some	Some	Some	Some	Some	Some	Some
Mening. B					Some	Some	Some	Some	Some	Some	Some	Some
Mpox							Some	Some	Some	Some	Some	Some
Pneumo. (PCV15)	Universal	Universal	Universal	Some	Some	Some	Some	Some	Some	Some	Some	Some
Pneumo. (PCV20)	Universal	Universal	Universal	Some	Some	Some	Some	Some	Some	Some	Some	Some
Pneumo. (PCV21)							Some	Some	Some	Some	Some	Some
Pneumo. (PPSV)			@ 2 years	Some	Some	Some	Some	Some	Some	Some	Some	Some
Polio	Universal	Universal	Universal	Universal	Universal	Universal	Some	Some	Some	Some	Some	Some
RSV antibody	Universal	Up to 7 months	Up to 19 months									
RSV vaccine						Some	Some	Some	Some	Some	Some	@ 75 years
Rotavirus	Universal	Up to 8 months										
Varicella			Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal	Universal
Zoster								Some	Some	Some	Some	Some

 Universal recommendation in the age group
 Recommendation for some individuals in the age group or shared clinical-decision making

Immunization recommendations for recipients of hematopoietic stem cell transplants may be found as indicated in chapter 2 of the [Pink Book](#).

Last revised May 21, 2026

Medicaid Covered Travel and Risk-Based Immunizations – as approved by the Oregon Immunization Program Medical Director. Click [here](#) for model protocols for clinical guidance.

	Travel	Pre-exposure risk	Post-exposure prophylaxis
Immune Globulin			
Hep. A (IG)			
Measles (IG)			
Hep. B (HBIG)			
Tetanus (TIG)			
Varicella (VZIG)			
Chikungunya	12 years +		
Cholera	2 years +		
Japanese Encephalitis	2 months +		
Mpox		18 years +	≥6 months of age
Rabies	All ages	All ages	
Tick-borne Encephalitis (TBE)	1 year +	Occupational only	
Typhoid	2 years +		
Yellow Fever	9 months +		