

## INSTRUCTIONS FOR HEALTHCARE PROFESSIONALS: Prescribing Naloxone

Naloxone is the antidote for an opioid overdose. It has been used for decades to reverse respiratory depression associated with toxic exposure to opioids. Naloxone is not a controlled substance and can be prescribed by anyone with a medical license. Take-home naloxone can be prescribed to patients at risk of an opioid overdose. Some reasons for prescribing naloxone are:

1. Receiving emergency medical care involving opioid intoxication or overdose
2. Suspected history of substance abuse or nonmedical opioid use
3. Starting methadone or buprenorphine for addiction
4. Higher-dose (>50 mg morphine equivalent/day) opioid prescription
5. Receiving any opioid prescription for pain plus:
  - a. Rotated from one opioid to another because of possible incomplete cross-tolerance
  - b. Smoking, COPD, emphysema, asthma, sleep apnea, respiratory infection, other respiratory illness
  - c. Renal dysfunction, hepatic disease, cardiac illness, HIV/AIDS
  - d. Known or suspected concurrent alcohol use
  - e. Concurrent benzodiazepine or other sedative prescription
  - f. Concurrent antidepressant prescription
6. Patients who may have difficulty accessing emergency medical services (distance, remoteness)
7. Voluntary request from patient or caregiver

Two naloxone formulations are available. Intra-muscular injection is cheaper but may be less attractive because it involves using a needle syringe. (IM syringes aren't widely used to inject controlled substances.) Intra-nasal (IN) spray is of comparable effectiveness, but may be more difficult to obtain at a pharmacy. Check with pharmacist to see whether IM or IN is more feasible.

### Billing for Clinical Encounter to Prescribe Naloxone

Most private health insurance, Medicare and Medicaid cover naloxone, but it varies by state.

Drug Abuse Screening Test—DAST-10			
These Questions Refer to the Past 12 Months			
1	Have you used drugs other than those required for medical reasons?	Yes	No
2	Do you abuse more than one drug at a time?	Yes	No
3	Are you unable to stop using drugs when you want to?	Yes	No
4	Have you ever had blackouts or flashbacks as a result of drug use?	Yes	No
5	Do you ever feel bad or guilty about your drug use?	Yes	No
6	Does your spouse (or parents) ever complain about your involvement with drugs?	Yes	No
7	Have you neglected your family because of your use of drugs?	Yes	No
8	Have you engaged in illegal activities in order to obtain drugs?	Yes	No
9	Have you ever experienced withdrawal symptoms (felt sick) when you stopped taking drugs?	Yes	No
10	Have you had medical problems as a result of your drug use (eg, memory loss, hepatitis, convulsions, bleeding)?	Yes	No

#### Screening, Brief Intervention & Referral to Treatment

SBIRT can be used to bill time for counseling a patient. Complete the DAST-10 and counsel patient on how to recognize overdose and how to administer naloxone, using the following sheets. Refer to drug treatment program if appropriate.

#### Billing codes

Commercial insurance: CPT 99408 (15 to 30 mins.)

Medicare: G0396 (15 to 30 mins.)

Medicaid: H0050 (per 15 mins.)

Guidelines for Interpretation of DAST-10		
Interpretation (Each "Yes" response = 1)		
Score	Degree of Problems Related to Drug Abuse	Suggested Action
0	No problems reported	Encouragement and education
1-2	Low level	Risky behavior – feedback and advice
3-5	Moderate level	Harmful behavior – feedback and counseling; possible referral for specialized assessment
6-8	Substantial level	Intensive assessment and referral

#### Pharmacist: Dispensing Naloxone

Many outpatient pharmacies do not stock naloxone but it can be easily ordered from major distributors. The nasal atomizer can be ordered from the manufacturer LMA (1-800-788-7999), but isn't usually covered by insurance (\$3 each). It may take 24 hours to set up an account with LMA, and the minimum order size is 25.