TABLE 7-8. Early Neonatal Deaths¹ by Weeks of Gestation and Weight Oregon Residents, Birth Cohort 2002

Birthweight (In Grams)	Total	Weeks of Gestation								
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+
Total	137	8	52	30	4	7	4	16	8	8
001-349	23	7	15	1	_	_	_	_	_	_
350-499	21	_	18	3	_	_	_	_	_	_
<500	44	7	33	4	_	_	_	_	_	_
500-749	34	1	15	16	2	_	_	_	_	_
750-999	12	_	2	8	1	_	_	1	_	_
1000-1249	6	_	_	2	1	2	1	_	_	–
1250-1499	1	_	_	_	_	1	_	_	_	–
1500-1999	7	_	_	_	_	3	1	1	_	2
2000-2499	3	_	_	_	_	_	1	1	1	–
<2500	107	8	50	30	4	6	3	3	1	2
2500-2999	8	_	_	_	_	_	1	5	1	1
3000-3499	11	_	_	_	_	_	_	7	1	3
3500-3999	5	_	_	_	_	1	_	1	3	_
4000-4499	3	_	_	_	-	_	_	_	1	2
4500+	1	_	_	_	_	_	_	_	1	_

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.

Quantity is zero.
Early neonatal death is defined as less than 7 days old.