| Risk Factor | Perinatal I ¹ | | | Perinatal II ² | | | Neonatal ³ | |
|-------------------------------|--------------------------|------|-------|---------------------------|------|-------|-----------------------|------|
| | No. | Rate | Ratio | No. | Rate | Ratio | No. | Rate |
| Total | 783 | 5.7 | 5.7 | 1,103 | 8.1 | 8.1 | 489 | 3.6 |
| Marital Status | | • | • | ., | •••• | •••• | | 0.0 |
| Married | 508 | 5.4 | 5.4 | 705 | 7.4 | 7.4 | 309 | 3.3 |
| Unmarried | 275 | 6.6 | 6.6 | 398 | 9.5 | 9.6 | 180 | 4.3 |
| Mother's Age | | | | | | | | |
| 10-14 | 2 | _ | _ | 2 | _ | _ | 1 | _ |
| 15-19 | 86 | 6.0 | 6.0 | 127 | 8.8 | 8.9 | 59 | 4.1 |
| 20-24 | 205 | 5.6 | 5.6 | 290 | 7.9 | 7.9 | 133 | 3.6 |
| 25-29 | 198 | 5.2 | 5.2 | 264 | 7.0 | 7.0 | 124 | 3.3 |
| 30-34 | 159 | 5.2 | 5.2 | 234 | 7.7 | 7.7 | 96 | 3.2 |
| 35-39 | 107 | 7.6 | 7.7 | 142 | 10.1 | 10.2 | 61 | 4.4 |
| 40-44 | 22 | 7.2 | 7.2 | 35 | 11.4 | 11.5 | 11 | 3.6 |
| 45+ | 2 | _ | _ | 7 | 36.1 | 37.0 | 2 | _ |
| Non-Hispanic | | | | | | | | |
| White | 562 | 5.6 | 5.6 | 814 | 8.1 | 8.1 | 362 | 3.6 |
| African American | 23 | 8.2 | 8.2 | 40 | 14.2 | 14.3 | 14 | 5.0 |
| American Indian | 14 | 6.8 | 6.9 | 23 | 11.2 | 11.3 | 9 | 4.4 |
| Asian ⁴ | 25 | 3.7 | 3.7 | 36 | 5.3 | 5.3 | 18 | 2.6 |
| Total Hispanic | 155 | 6.6 | 6.6 | 186 | 7.9 | 8.0 | 84 | 3.6 |
| Mother's Education | | | | | | | | |
| 8 th Grade or Less | 53 | 6.0 | 6.0 | 66 | 7.4 | 7.5 | 25 | 2.8 |
| Some High School | 125 | 6.7 | 6.7 | 172 | 9.2 | 9.2 | 83 | 4.5 |
| HS diploma/GED | 256 | 5.9 | 6.0 | 382 | 8.8 | 8.9 | 164 | 3.8 |
| More than High School | 291 | 4.6 | 4.6 | 404 | 6.3 | 6.3 | 185 | 2.9 |
| Start of Prenatal Care | | | | | | | | |
| 1 st Trimester | 582 | 5.2 | 5.2 | 838 | 7.5 | 7.6 | 381 | 3.4 |
| 2 nd Trimester | 124 | 6.2 | 6.2 | 161 | 8.0 | 8.0 | 69 | 3.4 |
| 3 rd Trimester | 16 | 4.0 | 4.0 | 26 | 6.6 | 6.6 | 10 | 2.5 |
| No Care | 61 | 43.0 | 44.0 | 78 | 54.4 | 56.3 | 29 | 20.9 |
| Tobacco Use | | | | | | | | |
| Yes | 125 | 7.1 | 7.2 | 194 | 11.0 | 11.1 | 74 | 4.2 |
| No | 616 | 5.2 | 5.2 | 865 | 7.3 | 7.4 | 387 | 3.3 |
| Alcohol Use | | | | | | | | |
| Yes | 16 | 9.5 | 9.6 | 23 | 13.6 | 13.7 | 7 | 4.2 |
| No | 722 | 5.5 | 5.5 | 1,032 | 7.8 | 7.8 | 453 | 3.4 |
| Multiple Birth | | | | | | | | |
| Yes | 95 | 24.0 | 24.2 | 132 | 33.2 | 33.6 | 84 | 21.4 |
| No | 688 | 5.2 | 5.2 | 971 | 7.3 | 7.3 | 405 | 3.1 |

TABLE 7-16. Perinatal Death Rates by Mother's Risk Factors,Oregon Residents, Birth Cohort 2000-2002

⁻ Quantity is zero or rate/ratio is based on fewer than five occurrences.

NOTE: Because of unreported items, the sum of all categories may not equal the total.

¹ Perinatal Definition I includes fetal deaths at 28 weeks of gestation or more and infant deaths of less than 7 days.

² Perinatal Definition II includes fetal deaths at 20 weeks of gestation or more and infant deaths of less than 28 days.

³ Neonatal deaths include infant deaths of less than 28 days.

⁴ Includes Chinese, Japanese, Filipino, and Other Asian & Pacific Islander.

Perinatal I and Perinatal II ratios and Neonatal rates are per 1,000 births. Perinatal I rates include all live biths and fetal deaths at 28 weeks gestation or more. Perinatal II rates include all live births and fetal deaths at 20 weeks of gestation or more.