

**TABLE 7-9. Late Neonatal Deaths by Weeks of Gestation and Weight,
Oregon Residents, Birth Cohort 2003**

Birthweight (In Grams)	Total	Weeks of Gestation								
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+
Total	41	–	1	7	7	3	–	13	7	3
001-349	–	–	–	–	–	–	–	–	–	–
350-499	2	–	–	1	1	–	–	–	–	–
<500	2	–	–	1	1	–	–	–	–	–
500-749	3	–	–	2	1	–	–	–	–	–
750-999	6	–	1	3	2	–	–	–	–	–
1000-1249	3	–	–	1	1	1	–	–	–	–
1250-1499	3	–	–	–	2	1	–	–	–	–
1500-1999	2	–	–	–	–	1	–	1	–	–
2000-2499	4	–	–	–	–	–	–	4	–	–
<2500	23	–	1	7	7	3	–	5	–	–
2500-2999	2	–	–	–	–	–	–	2	–	–
3000-3499	9	–	–	–	–	–	–	3	5	1
3500-3999	4	–	–	–	–	–	–	3	1	–
4000-4499	–	–	–	–	–	–	–	–	–	–
4500-4999	2	–	–	–	–	–	–	–	1	1

– Quantity is zero.

Late neonatal death is defined as death at 7 to 27 days old.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used.